

CL

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 April 2001 (19.04.2001)


PCT

(10) International Publication Number
WO 01/27108 A1

- (51) International Patent Classification⁷: C07D 413/12, A61K 31/55, C07D 409/12, 401/12, 409/14, 403/12, 407/12, 243/10
- (74) Agent: LARSEN, Scott, K.; Du Pont Pharmaceuticals Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).
- (21) International Application Number: PCT/US00/27666
- (81) Designated States (*national*): AU, BR, CA, CN, CZ, EE, HU, IL, IN, JP, KR, LT, LV, MX, NO, NZ, PL, RO, SG, SI, SK, TR, UA, VN, ZA.
- (22) International Filing Date: 7 October 2000 (07.10.2000)
- (25) Filing Language: English
- (84) Designated States (*regional*): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- (26) Publication Language: English
- (30) Priority Data: 60/158,565 8 October 1999 (08.10.1999) US
- Published: — With international search report.
- (71) Applicant: DU PONT PHARMACEUTICALS COMPANY [US/US]; Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 (US).
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.
- (72) Inventors: THOMPSON, Lorin, Andrew; 600 Silverside Road, Wilmington, DE 19707 (US). HAN, Amy, Qi; 177 Thompson Drive, Hockessin, DE 19707 (US).



(54) Title: AMINO LACTAM SULFONAMIDES AS INHIBITORS OF A β PROTEIN PRODUCTION



WO 01/27108 A1

(57) Abstract: This invention relates to novel lactams having Formula (I), to their pharmaceutical compositions and to their methods of use. These novel compounds inhibit the processing of amyloid precursor protein and, more specifically, inhibit the production of A β -peptide, thereby acting to prevent the formation of neurological deposits of amyloid protein. More particularly, the present invention relates to the treatment of neurological disorders related to β -amyloid production such as Alzheimer's disease and Down's Syndrome.

TITLEAMINO LACTAM SULFONAMIDES AS INHIBITORS OF A β PROTEIN
PRODUCTION5 FIELD OF THE INVENTION

This invention relates to novel lactams having drug and bio-affecting properties, their pharmaceutical compositions and methods of use. These novel compounds inhibit the processing of amyloid precursor protein and, more specifically, inhibit the production of A β -peptide, thereby acting to prevent the formation of neurological deposits of amyloid protein. More particularly, the present invention relates to the treatment of neurological disorders related to β -amyloid production such as Alzheimer's disease and Down's Syndrome.

BACKGROUND OF THE INVENTION

Alzheimer's disease (AD) is a degenerative brain disorder characterized clinically by progressive loss of memory, temporal and local orientation, cognition, reasoning, judgment and emotionally stability. AD is a common cause of progressive dementia in humans and is one of the major causes of death in the United States. AD has been observed in all races and ethnic groups worldwide, and is a major present and future health problem. No treatment that effectively prevents AD or reverses the clinical symptoms and underlying pathophysiology is currently available (for review, Dennis J. Selkoe; Cell Biology of the amyloid (beta)-protein precursor and the mechanism of Alzheimer's disease, Annu Rev Cell Biol, 1994, 10: 373-403).

Histopathological examination of brain tissue derived upon autopsy or from neurosurgical specimens in effected individuals revealed the occurrence of amyloid plaques and neurofibrillar tangles in the cerebral cortex of such patients. Similar alterations were observed in patients with Trisomy 21 (Down's syndrome), and hereditary cerebral

hemorrhage with amyloidosis of the Dutch-type.

Neurofibrillar tangles are nonmembrane-bound bundles of abnormal proteinaceous filaments and biochemical and immunochemical studies led to the conclusion that their principle protein subunit is an altered phosphorylated form of the tau protein (reviewed in Selkoe, 1994).

Biochemical and immunological studies revealed that the dominant proteinaceous component of the amyloid plaque is an approximately 4.2 kilodalton (kD) protein of about 39 to 43 amino acids. This protein was designated A β , β -amyloid peptide, and sometimes β /A4; referred to herein as A β . In addition to its deposition in amyloid plaques, A β is also found in the walls of meningeal and parenchymal arterioles, small arteries, capillaries, and sometimes, venules. A β was first purified and a partial amino acid reported in 1984 (Glenner and Wong, Biochem. Biophys. Res. Commun. 120: 885-890). The isolation and sequence data for the first 28 amino acids are described in U.S. Pat. No 4,666,829.

Compelling evidence accumulated during the last decade revealed that A β is an internal polypeptide derived from a type 1 integral membrane protein, termed β amyloid precursor protein (APP). β APP is normally produced by many cells both in vivo and in cultured cells, derived from various animals and humans. A β is derived from cleavage of β APP by as yet unknown enzyme (protease) system(s), collectively termed secretases.

The existence of at least four proteolytic activities has been postulated. They include β secretase(s), generating the N-terminus of A β , α secretase(s) cleaving around the 16/17 peptide bond in A β , and γ secretases, generating C-terminal A β fragments ending at position 38, 39, 40, 42, and 43 or generating C-terminal extended precursors which are subsequently truncated to the above polypeptides.

Several lines of evidence suggest that abnormal accumulation of A β plays a key role in the pathogenesis of AD. Firstly, A β is the major protein found in amyloid plaques. Secondly, A β is neurotoxic and may be causally related to neuronal death observed in AD patients. Thirdly, missense DNA mutations at position 717 in the 770 isoform of β APP can be found in effected members but not unaffected members of several families with a genetically determined (familiar) form of AD. In addition, several other β APP mutations have been described in familiar forms of AD. Fourthly, similar neuropathological changes have been observed in transgenic animals overexpressing mutant forms of human β APP. Fifthly, individuals with Down's syndrome have an increased gene dosage of β APP and develop early-onset AD. Taken together, these observations strongly suggest that A β depositions may be causally related to the AD.

It is hypothesized that inhibiting the production of A β will prevent and reduce neurological degeneration, by controlling the formation of amyloid plaques, reducing neurotoxicity and, generally, mediating the pathology associated with A β production. One method of treatment methods would therefore be based on drugs that inhibit the formation of A β in vivo.

Methods of treatment could target the formation of A β through the enzymes involved in the proteolytic processing of β amyloid precursor protein. Compounds that inhibit β or γ secretase activity, either directly or indirectly, could control the production of A β . Advantageously, compounds that specifically target γ secretases, could control the production of A β . Such inhibition of β or γ secretases could thereby reduce production of A β , which, thereby, could reduce or prevent the neurological disorders associated with A β protein.


SUMMARY OF THE INVENTION

One object of the present invention is to provide novel compounds which are useful as inhibitors of the production of A β protein or pharmaceutically acceptable salts or prodrugs thereof.

It is another object of the present invention to provide pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt or prodrug form thereof.

It is another object of the present invention to provide a method for treating degenerative neurological disorders comprising administering to a host in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt or prodrug form thereof.

These and other objects, which will become apparent during the following detailed description, have been achieved by the inventors' discovery that compounds of Formula (I):




(I)

or pharmaceutically acceptable salt or prodrug forms thereof, wherein Q, R², R³, R⁵, R^{5a}, R⁶, B, W, X, Y, and Z are defined below, are effective inhibitors of the production of A β .

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Thus, in one embodiment, the present invention provides a novel compound of Formula (I):



or a pharmaceutically acceptable salt or prodrug thereof,
 5 wherein:

Q is C₁-C₆ alkyl substituted with 0-3 R^{1a};
 C₂-C₆ alkenyl substituted with 0-3 R^{1a};
 C₂-C₆ alkynyl substituted with 0-3 R^{1a};
 10 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1b};
 C₃-C₁₀ carbocycle substituted with 0-3 R^{1b};
 C₆-C₁₀ aryl substituted with 0-3 R^{1b}; or
 5 to 10 membered heterocycle substituted with 0-3 R^{1b};

15 R^{1a}, at each occurrence, is independently selected from H,
 R^{1b}, Cl, F, Br, I, OR¹⁴, CN, NO₂, =O, NR¹⁹R²⁰, CF₃,
 C₁-C₄ alkyl; C₂-C₄ alkenyl; C₂-C₄ alkynyl; C₁-C₄
 haloalkyl; C₁-C₄ haloalkoxy;
 C₃-C₁₀ carbocycle substituted with 0-3 R^{1b};
 20 C₆-C₁₀ aryl substituted with 0-3 R^{1b}; and
 5 to 10 membered heterocycle substituted with 0-3 R^{1b};

25 R^{1b}, at each occurrence, is independently selected from H,
 OR¹⁴, Cl, F, Br, I, CN, NO₂, =O, NR¹⁹R²⁰, CF₃,
 C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;
 -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d}, -S(=O)-R^{1d}, -S(=O)₂-R^{1d},
 -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d},
 -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d}, -NR^{19b}S(=O)-R^{1d},
 30 -S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, -OC(=O)-R^{1d};

C₁-C₆ alkyl substituted with 0-3 R^{1c};

- C₂-C₆ alkenyl substituted with 0-2 R^{1c};
- C₂-C₆ alkynyl substituted with 0-2 R^{1c};
- C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
- C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};
- 5 C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
- 5 to 10 membered heterocycle substituted with 0-3 R^{1f};

- R^{1c}, at each occurrence, is independently selected from H,
 OR¹⁴, Cl, F, Br, I, CN, NO₂, NR¹⁹R²⁰, CF₃,
- 10 C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;
 -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d}, -S(=O)-R^{1d}, -S(=O)₂-R^{1d},
 -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d},
 -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d}, -NR^{19b}S(=O)-R^{1d},
 -S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, -OC(=O)-R^{1d};

- 15 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
- C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};
- C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
- 5 to 10 membered heterocycle substituted with 0-3 R^{1f};

- 20 R^{1d}, at each occurrence, is independently selected from H,
 C₁-C₆ alkyl substituted with 0-3 R^{1e};
- C₂-C₆ alkenyl substituted with 0-2 R^{1e};
- C₂-C₆ alkynyl substituted with 0-2 R^{1e};
- 25 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
- C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};
- C₆-C₁₀ aryl substituted with 0-3 R^{1f};
- 5 to 10 membered heterocycle substituted with 0-3 R^{1f};

- 30 R^{1e}, at each occurrence, is independently selected from H,
 OR¹⁴, Cl, F, Br, I, CN, NO₂, =O, NR¹⁹R²⁰, CF₃, C₁-C₄
 alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;
 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
- C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};

C₆-C₁₀ aryl substituted with 0-3 R^{1f};
5 to 10 membered heterocycle substituted with 0-3 R^{1f};

5 R^{1f}, at each occurrence, is independently selected
from H, OR¹⁴, Cl, F, Br, I, CN, NO₂, =O, NR¹⁹R²⁰, CF₃,
C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, and C₁-C₄
haloalkoxy;

10 R² is H, C₁-C₆ alkyl, (C₁-C₆ alkoxy) C₁-C₃ alkyl,
C₃-C₆ carbocycle, C₆-C₁₀ aryl, (C₃-C₆
carbocycle)methyl, (C₆-C₁₀ aryl)methyl, (C₆-C₁₀
aryl)ethyl, or 5 to 10 membered heterocycle;

15 R⁵ is H, OH, OR¹⁴;
C₁-C₆ alkyl substituted with 0-3 R^{5b};
C₁-C₆ alkoxy substituted with 0-3 R^{5b};
C₂-C₆ alkenyl substituted with 0-3 R^{5b};
C₂-C₆ alkynyl substituted with 0-3 R^{5b};
20 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{5c};
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or
5 to 10 membered heterocycle substituted with 0-3R^{5c};

25 R^{5a} is H, OH, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₂-C₄ alkenyl, or
C₂-C₄ alkenyloxy;

alternatively, R⁵ and R^{5a} may be combined to form =R⁵;

30 alternatively, R⁵ and R^{5a} may be combined to form a 3-7
membered cycloalkyl ring substituted with 0-3 R^{5c};
optionally the cycloalkyl ring formed by combining R⁵
and R^{5a} may be benzo fused, wherein the benzo fused
ring may be substituted with 0-3 R^{5c};

35 R^{5b}, at each occurrence, is independently selected from:

- H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
NR¹⁵R¹⁶;
C₃-C₁₀ cycloalkyl substituted with 0-3 R^{5c};
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
5 C₆-C₁₀ aryl substituted with 0-3 R^{5c}; and
5 to 10 membered heterocycle substituted with 0-3 R^{5c};
- R^{5c}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
10 S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ halothioalkoxy;
- R⁶ is H;
C₁-C₆ alkyl substituted with 0-3 R^{6a};
15 C₃-C₁₀ carbocycle substituted with 0-3 R^{6b}; or
C₆-C₁₀ aryl substituted with 0-3 R^{6b};
- R^{6a}, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
20 aryl or CF₃;
- R^{6b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, C₁-C₆ alkyl,
C₁-C₄ alkoxy, C₁-C₄ haloalkyl, and C₁-C₄ haloalkoxy;
25
- Ring B is a 5 to 10 membered lactam,
wherein the lactam is saturated, partially saturated
or unsaturated;
wherein each additional lactam carbon is substituted
30 with 0-2 R¹¹; and,
optionally, the lactam contains a heteroatom selected
from -N-, -NH-, -N(R¹⁰)-, -O-, -S-, -S(=O)-, and
-S(=O)₂-;
- 35 additionally, two R¹¹ substituents on adjacent atoms may be
combined to form C₃-C₆ carbocycle fused radical, a

benzo fused radical, or a 5 to 6 membered heteroaryl
fused radical,
wherein said 5 to 6 membered heteroaryl fused radical
comprises 1-2 heteroatoms selected from N, O, and S;
5 wherein said benzo fused radical or 5 to 6 membered
heteroaryl fused radical is substituted with 0-3 R¹³;

R¹⁰ is H, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹,
S(=O)₂R¹⁷;

10 C₁-C₆ alkyl substituted with 0-2 R^{10a};
C₆-C₁₀ aryl substituted with 0-4 R^{10b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{10b}; or
5 to 10 membered heterocycle optionally substituted
with 0-3 R^{10b};

15 R^{10a}, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
CF₃, or aryl substituted with 0-4 R^{10b};

20 R^{10b}, at each occurrence, is independently selected from H,
OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂,
NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆
alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy,
and C₁-C₄ halothioalkoxy;

25 R¹¹, at each occurrence, is independently selected from
C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹,
C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹,
CF₃;

30 C₁-C₆ alkyl substituted with 0-1 R^{11a};
C₆-C₁₀ aryl substituted with 0-3 R^{11b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
5 to 10 membered heterocycle substituted with 0-3
R^{11b};

35

R^{11a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, CF₃, or phenyl substituted with 0-3 R^{11b};

5 R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ halothioalkoxy;

10 W is -(CR⁸R^{8a})_p-;

p is 0, 1, 2, 3, or 4;

15 R⁸ and R^{8a}, at each occurrence, are independently selected from H, F, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl and C₃-C₈ cycloalkyl;

X is a bond;

C₆-C₁₀ aryl substituted with 0-3 R^{Xb};

20 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{Xb};

C₃-C₁₀ carbocycle substituted with 0-3 R^{Xb}; or

5 to 10 membered heterocycle substituted with 0-2 R^{Xb};

25 R^{Xb}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ halothioalkoxy;

Y is a bond or -(CR⁹R^{9a})_t-V-(CR⁹R^{9a})_u-;

30

t is 0, 1, 2, or 3;

u is 0, 1, 2, or 3;

R⁹ and R^{9a}, at each occurrence, are independently selected from H, F, C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

5 V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -N(R¹⁹)-, -C(=O)NR^{19b}-, -NR^{19b}C(=O)-, -NR^{19b}S(=O)₂-, -S(=O)₂NR^{19b}-, -NR^{19b}S(=O)-, -S(=O)NR^{19b}-, -C(=O)O-, or -OC(=O)-;

Z is H;

10 C₁-C₈ alkyl substituted with 0-2 R¹²;
C₂-C₄ alkenyl substituted with 0-2 R¹²;
C₂-C₄ alkynyl substituted with 0-2 R¹²;
C₆-C₁₀ aryl substituted with 0-4 R^{12b};
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or
15 5 to 10 membered heterocycle substituted with 0-3 R^{12b};

R¹² is C₆-C₁₀ aryl substituted with 0-4 R^{12b};
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or
20 5 to 10 membered heterocycle substituted with 0-3 R^{12b};

R^{12b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
25 S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ halothioalkoxy;

R¹³, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂,
30 NR¹⁵R¹⁶, or CF₃;

R¹⁴, at each occurrence, is independently selected from H, phenyl, benzyl, C₁-C₆ alkyl, or C₂-C₆ alkoxyalkyl;

R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

5 R¹⁶, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

10 R¹⁷ is H, aryl, aryl-CH₂-, C₁-C₆ alkyl, or C₂-C₆ alkoxyalkyl;

R¹⁸, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

15


R¹⁹, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

20 R^{19b}, at each occurrence, is independently selected from H, C₁-C₄ alkyl, phenyl, benzyl, and phenethyl; and

R²⁰ and R²¹, at each occurrence, are independently selected from H, C₁-C₄ alkyl, aryl, and aryl(C₁-C₂ alkyl)-.

25

[1] In another embodiment, the present invention provides a novel compound of Formula (I):



30

(I)

or a pharmaceutically acceptable salt or prodrug thereof, wherein:

- Q is C₁-C₆ alkyl substituted with 0-3 R^{1a};
 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1b};
 C₃-C₁₀ carbocycle substituted with 0-3 R^{1b};
 C₆-C₁₀ aryl substituted with 0-3 R^{1b}; or
 5 5 to 10 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered
 heterocycle is substituted with 0-3 R^{1b};
- 10 R^{1a}, at each occurrence, is independently selected from H,
 R^{1b}, Cl, F, Br, I, OR¹⁴, CN, NO₂, =O, NR¹⁹R²⁰, CF₃,
 C₁-C₄ alkyl; C₂-C₄ alkenyl; C₂-C₄ alkynyl; C₁-C₄
 haloalkyl; C₁-C₄ haloalkoxy;
- 15 C₃-C₁₀ carbocycle substituted with 0-3 R^{1b};
 C₆-C₁₀ aryl substituted with 0-3 R^{1b}; and
 5 to 10 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered
 20 heterocycle is substituted with 0-3 R^{1b};
- R^{1b}, at each occurrence, is independently selected from H,
 OR¹⁴, Cl, F, Br, I, CN, NO₂, =O, NR¹⁹R²⁰, CF₃, OCF₃,
 C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;
- 25 -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d}, -S(=O)-R^{1d}, -S(=O)₂-R^{1d},
 -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d},
 -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d}, -NR^{19b}S(=O)-R^{1d},
 -S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, -OC(=O)-R^{1d};
- 30 C₁-C₆ alkyl substituted with 0-3 R^{1c};
 C₂-C₆ alkenyl substituted with 0-2 R^{1c};
 C₂-C₆ alkynyl substituted with 0-2 R^{1c};
 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};

- C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};
C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
5 to 14 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 14 membered
heterocycle is substituted with 0-3 R^{1f};
- R^{1c}, at each occurrence, is independently selected from H,
OR¹⁴, Cl, F, Br, I, CN, NO₂, NR¹⁹R²⁰, CF₃,
C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;
-C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d}, -S(=O)-R^{1d}, -S(=O)₂-R^{1d},
-N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d},
-NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d}, -NR^{19b}S(=O)-R^{1d},
-S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, -OC(=O)-R^{1d};
- C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};
C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{1f};
- R^{1d}, at each occurrence, is independently selected from H,
C₁-C₆ alkyl substituted with 0-3 R^{1e};
C₂-C₆ alkenyl substituted with 0-2 R^{1e};
C₂-C₆ alkynyl substituted with 0-2 R^{1e};
C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};
C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{1f};

R^{1e}, at each occurrence, is independently selected from H,
5 OR¹⁴, Cl, F, Br, I, CN, NO₂, =O, NR¹⁹R²⁰, CF₃, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;
C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};
10 C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{1f};

15 R^{1f}, at each occurrence, is independently selected from H, OR¹⁴, SR¹⁴, Cl, F, Br, I, CN, NO₂, =O, NR¹⁹R²⁰, CF₃, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

20 R² is H, C₁-C₆ alkyl, (C₁-C₆ alkoxy) C₁-C₃ alkyl, C₃-C₆ carbocycle, C₆-C₁₀ aryl, (C₃-C₆ carbocycle)methyl, (C₆-C₁₀ aryl)methyl, (C₆-C₁₀ aryl)ethyl, or 5 to 10 membered heterocycle;

25 R⁵ is H, OR¹⁴;
C₁-C₆ alkyl substituted with 0-3 R^{5b};
C₁-C₆ alkoxy substituted with 0-3 R^{5b};
C₂-C₆ alkenyl substituted with 0-3 R^{5b};
30 C₂-C₆ alkynyl substituted with 0-3 R^{5b};
C₃-C₁₀ cycloalkyl substituted with 0-3 R^{5c};
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or

- 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};
- 5 R^{5a} is H or C₁-C₄ alkyl;
- R^{5b}, at each occurrence, is independently selected from:
H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
10 NR¹⁵R¹⁶; acetyl, -S(C₁-C₄ alkyl), -S(=O)(C₁-C₄ alkyl), -S(=O)₂(C₁-C₄ alkyl);
C₃-C₁₀ cycloalkyl substituted with 0-3 R^{5c};
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
C₆-C₁₀ aryl substituted with 0-3 R^{5c}; and
15 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};
- 20 R^{5c}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;
- 25 R⁶ is H;
C₁-C₆ alkyl substituted with 0-3 R^{6a};
C₃-C₁₀ carbocycle substituted with 0-3 R^{6b}; or
C₆-C₁₀ aryl substituted with 0-3 R^{6b};
- 30 R^{6a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, aryl or CF₃;

- R^{6b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, and C₁-C₄ haloalkoxy;
- 5 Ring B is a 6, 7, or 8 membered lactam,
 wherein the lactam is saturated, partially saturated
 or unsaturated;
 wherein each additional lactam carbon is substituted
 with 0-2 R¹¹; and,
10 optionally, the lactam contains a heteroatom selected
 from -N=, -NH-, -N(R¹⁰)-, -O-, -S-, -S(=O)-, and
 -S(=O)₂-;
- additionally, two R¹¹ substituents on adjacent atoms may be
15 combined to form C₃-C₆ carbocycle fused radical, a
 benzo fused radical, or a 5 to 6 membered heteroaryl
 fused radical,
 wherein said 5 to 6 membered heteroaryl fused radical
 comprises 1-2 heteroatoms selected from N, O, and S;
20 wherein said benzo fused radical or 5 to 6 membered
 heteroaryl fused radical is substituted with 0-3
 R¹³;
- R¹⁰ is H, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹,
25 S(=O)₂R¹⁷;
 C₁-C₆ alkyl substituted with 0-2 R^{10a};
 C₆-C₁₀ aryl substituted with 0-4 R^{10b};
 C₃-C₁₀ carbocycle substituted with 0-3 R^{10b}; or
 5 to 10 membered heterocycle containing 1 to 4
30 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered
 heterocycle is substituted with 0-3 R^{10b};

R^{10a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, CF₃, or aryl substituted with 0-4 R^{10b};

5 R^{10b}, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

10

R¹¹, at each occurrence, is independently selected from H, C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, CF₃;

15

C₁-C₆ alkyl substituted with 0-1 R^{11a};
C₆-C₁₀ aryl substituted with 0-3 R^{11b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
20 sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{11b};

R^{11a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
25 CF₃, or phenyl substituted with 0-3 R^{11b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
30 is substituted with 0-3 R^{11b};

R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,

S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

W is -(CR⁸R^{8a})_p-;

5

p is 0, 1, 2, 3, or 4;

R⁸ and R^{8a}, at each occurrence, are independently selected from H, F, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl and C₃-C₈ cycloalkyl;

10

X is a bond;

C₆-C₁₀ aryl substituted with 0-3 R^{Xb};

C₃-C₁₀ cycloalkyl substituted with 0-3 R^{Xb};

15

C₃-C₁₀ carbocycle substituted with 0-3 R^{Xb}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered

V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -N(R¹⁹)-, -C(=O)NR^{19b}-, -NR^{19b}C(=O)-, -NR^{19b}S(=O)₂-, -S(=O)₂NR^{19b}-, -NR^{19b}S(=O)-, -S(=O)NR^{19b}-, -C(=O)O-, or -OC(=O)-;

5

Z is H;

C₁-C₈ alkyl substituted with 0-2 R¹²;

C₂-C₆ alkenyl substituted with 0-2 R¹²;

C₂-C₆ alkynyl substituted with 0-2 R¹²;

10 C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered

15 heterocycle is substituted with 0-3 R^{12b};

R¹² is C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or

20 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

25 R^{12b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

30 R¹³, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, or CF₃;

R¹⁴, at each occurrence, is independently selected from H, phenyl, benzyl, C₁-C₆ alkyl, or C₂-C₆ alkoxyalkyl;

5 R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

10 R¹⁶, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

alternatively, -NR¹⁵R¹⁶ may be a heterocyclic ring selected from the group piperidinyl, morpholinyl,
15 thiomorpholinyl, pyrrolidinyl, homopiperidinyl, piperazinyl, and N-methylpiperizinyl;

R¹⁷ is H, aryl, aryl-CH₂-, C₁-C₆ alkyl, or C₂-C₆ alkoxyalkyl;

20


R¹⁸, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

25 R¹⁹, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

30 R^{19b}, at each occurrence, is independently selected from H, C₁-C₄ alkyl, phenyl, benzyl, and phenethyl; and

R²⁰ and R²¹, at each occurrence, are independently selected from H, C₁-C₄ alkyl, aryl, and aryl(C₁-C₂ alkyl)-;

provided that when Q is phenyl, 3-F-phenyl, benzyl, propyl, or octyl; R² is H; R⁵ is methyl, and -WXYZ is methyl, then Ring B is not



5

; wherein R¹³ is H.

[2] In an alternative embodiment of the present invention of of Formula (I),

10

Q is C₁-C₆ alkyl substituted with 0-3 R^{1a};

C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{1b};

C₆-C₁₀ aryl substituted with 0-3 R^{1b}; or

15

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{1b};

20 R^{1a}, at each occurrence, is independently selected from H, R^{1b}, Cl, F, Br, NR¹⁹R²⁰, CF₃, C₁-C₄ alkyl, C₁-C₄ haloalkyl;

C₃-C₁₀ carbocycle substituted with 0-3 R^{1b};

C₆-C₁₀ aryl substituted with 0-3 R^{1b}; and

25

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{1b};

30 R^{1b}, at each occurrence, is independently selected from H, OR¹⁴, Cl, F, Br, NO₂, NR¹⁹R²⁰, CF₃, OCF₃,

- C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;
 -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d}, -S(=O)-R^{1d}, -S(=O)₂-R^{1d},
 -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d},
 -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d}, -NR^{19b}S(=O)-R^{1d},
 5 -S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, -OC(=O)-R^{1d};
 C₁-C₆ alkyl substituted with 0-3 R^{1c};
 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
 C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
 5 to 14 membered heterocycle containing 1 to 4
 10 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 14 membered
 heterocycle is substituted with 0-3 R^{1f};

 R^{1c}, at each occurrence, is independently selected from
 15 H, OR¹⁴, Cl, F, Br, NO₂, NR¹⁹R²⁰, CF₃, C₁-C₄ alkoxy,
 C₁-C₄ haloalkoxy; -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d},
 -S(=O)-R^{1d}, -S(=O)₂-R^{1d}, -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d},
 -NR^{19b}C(=O)-R^{1d}, -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d},
 -NR^{19b}S(=O)-R^{1d}, -S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d},
 20 -OC(=O)-R^{1d};

 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
 C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};
 C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
 25 5 to 10 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered
 heterocycle is substituted with 0-3 R^{1f};

 30 R^{1d}, at each occurrence, is independently selected from H,
 C₁-C₆ alkyl, and C₆-C₁₀ aryl substituted with 0-3 R^{1f};

 R^{1f}, at each occurrence, is independently selected from

H, OR¹⁴, SR¹⁴, Cl, F, Br, I, CN, NO₂, =O, NR¹⁹R²⁰,
CF₃, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

5 R² is H or C₁-C₆ alkyl;

R⁵ is H, OR¹⁴;

C₁-C₆ alkyl substituted with 0-3 R^{5b};

C₁-C₆ alkoxy substituted with 0-3 R^{5b};

10 C₂-C₆ alkenyl substituted with 0-3 R^{5b}; or

C₂-C₆ alkynyl substituted with 0-3 R^{5b};

R^{5a} is H or C₁-C₄ alkyl;

15 R^{5b}, at each occurrence, is independently selected from:

H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
NR¹⁵R¹⁶; acetyl, S(C₁-C₄ alkyl), S(=O)(C₁-C₄
alkyl), S(=O)₂(C₁-C₄ alkyl),

C₃-C₁₀ cycloalkyl substituted with 0-3 R^{5c};

20 C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};

C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered

25 heterocycle is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
30 haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁶ is H, methyl, or ethyl;

- Ring B is a seven membered lactam,
wherein the lactam is saturated, partially saturated
or unsaturated;
wherein each additional lactam carbon is substituted
5 with 0-2 R¹¹; and,
optionally, the lactam contains a heteroatom selected
from -N=, -NH-, -N(R¹⁰)-, -O-, -S-, -S(=O)-, and
-S(=O)₂-;
- 10 additionally, two R¹¹ substituents on adjacent atoms may be
combined to form C₃-C₆ carbocycle fused radical, a
benzo fused radical, or a 5 to 6 membered heteroaryl
fused radical,
wherein said 5 to 6 membered heteroaryl fused radical
15 comprises 1-2 heteroatoms selected from N, O, and S;
wherein said benzo fused radical or 5 to 6 membered
heteroaryl fused radical is substituted with 0-3
R¹³;
- 20 R¹⁰ is H, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹,
S(=O)₂R¹⁷;
C₁-C₆ alkyl substituted with 0-2 R^{10a};
C₆-C₁₀ aryl substituted with 0-4 R^{10b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{10b}; or
25 5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{10b};
- 30 R^{10a}, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
CF₃, or aryl substituted with 0-4 R^{10b};

R^{10b}, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R¹¹, at each occurrence, is independently selected from H, C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, CF₃;
C₁-C₆ alkyl substituted with 0-1 R^{11a};
C₆-C₁₀ aryl substituted with 0-3 R^{11b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{11b};

R^{11a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, CF₃, or phenyl substituted with 0-3 R^{11b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{11b};

R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

W is -(CR⁸R^{8a})_p-;

p is 0, 1, or 2;

5 R^8 and R^{8a} , at each occurrence, are independently selected from H, F, methyl, and ethyl;

X is a bond;

phenyl substituted with 0-3 R^{Xb} ;

C₃-C₆ cycloalkyl substituted with 0-3 R^{Xb} ; or

10 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-2 R^{Xb} ;

15 R^{Xb} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

20 Y is a bond or $-(CR^9R^{9a})_t-V-(CR^9R^{9a})_u-$;

t is 0, 1, or 2;

u is 0, 1, or 2;

25

R^9 and R^{9a} , at each occurrence, are independently selected from H, F, methyl, and ethyl;

V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-,

30 -N(R¹⁹)-, -NHC(=O)-, or -C(=O)NH-;

Z is H;

C₁-C₆ alkyl substituted with 0-2 R^{12} ;

C₂-C₆ alkenyl substituted with 0-2 R^{12} ;

35 C₂-C₆ alkynyl substituted with 0-2 R^{12} ;

- phenyl substituted with 0-4 R^{12b};
C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b};
- R¹² is phenyl substituted with 0-4 R^{12b};
C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b};
- R^{12b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;
- R¹³, at each occurrence, is independently selected from
H, OH, C₁-C₄ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
NO₂, NR¹⁵R¹⁶, and CF₃;
- R¹⁴, at each occurrence, is independently selected from H,
phenyl, benzyl, C₁-C₆ alkyl, and C₂-C₆ alkoxyalkyl;
- R¹⁵, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆
alkyl) and -S(=O)₂-(C₁-C₆ alkyl);
- R¹⁶, at each occurrence, is independently selected from H,
OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-
(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

alternatively, $-NR^{15}R^{16}$ may be a heterocyclic ring selected from the group piperidinyl, morpholinyl, thiomorpholinyl, pyrrolidinyl, homopiperidinyl, piperazinyl, and N-methylpiperizinyll;

5


R^{17} is H, aryl, aryl- CH_2 -, C_1 - C_6 alkyl, or C_2 - C_6 alkoxyalkyl;

10 R^{18} , at each occurrence, is independently selected from H, C_1 - C_6 alkyl, phenyl, benzyl, phenethyl, $-C(=O)-(C_1-C_6$ alkyl) and $-S(=O)_2-(C_1-C_6$ alkyl);

15 R^{19} , at each occurrence, is independently selected from H, OH, C_1 - C_6 alkyl, phenyl, benzyl, phenethyl, $-C(=O)-(C_1-C_6$ alkyl) and $-S(=O)_2-(C_1-C_6$ alkyl); and

R^{20} is H or C_1 - C_4 ;

20 provided that when Q is phenyl, 3-F-phenyl, benzyl, propyl, or octyl; R^2 is H; R^5 is methyl, and -WXYZ is methyl, then Ring B is not




25

[3] In another alterantive embodiment of the present invention of of Formula (I),

Ring B is selected from:

30



wherein each benzo fused radical is substituted with 0-3
R¹³.

[4] In another alternative embodiment of the present
invention of of Formula (I),

- Q is C₁-C₆ alkyl substituted with 0-3 R^{1a};
C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1b};
C₆-C₁₀ aryl substituted with 0-3 R^{1b}; or

- 5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{1b};
- 5 R^{1a}, at each occurrence, is independently selected from
H, Cl, F, Br, NR¹⁹R²⁰, CF₃, C₁-C₄ alkyl; C₁-C₄
haloalkyl;
C₆-C₁₀ aryl substituted with 0-3 R^{1b}; and
- 10 5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{1b};
- 15 R^{1b}, at each occurrence, is independently selected from
H, OR¹⁴, Cl, F, Br, NO₂, NR¹⁹R²⁰, CF₃, OCF₃,
C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;
-C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d}, -S(=O)-R^{1d}, -S(=O)₂-R^{1d},
-N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d},
20 -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d}, -NR^{19b}S(=O)-R^{1d},
-S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, -OC(=O)-R^{1d};
- C₁-C₆ alkyl substituted with 0-3 R^{1c};
C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
- 25 C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
5 to 14 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 14 membered
heterocycle is substituted with 0-3 R^{1f};
- 30 R^{1c}, at each occurrence, is independently selected from H,
OR¹⁴, Cl, F, Br, NO₂, NR¹⁹R²⁰, CF₃, C₁-C₄ alkoxy,
C₁-C₄ haloalkoxy; -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d},

-S(=O)-R^{1d}, -S(=O)₂-R^{1d}, -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d},
 -NR^{19b}C(=O)-R^{1d}, -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d},
 -NR^{19b}S(=O)-R^{1d}, -S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d},
 -OC(=O)-R^{1d};

5

C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};

C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};

C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and

5 to 10 membered heterocycle containing 1 to 4
 10 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered
 heterocycle is substituted with 0-3 R^{1f};

R^{1d}, at each occurrence, is independently selected from
 15 H, C₁-C₆ alkyl, and
 C₆-C₁₀ aryl substituted with 0-3 R^{1f};

R^{1f}, at each occurrence, is independently selected from
 H, Cl, F, Br, NO₂, CF₃, C₁-C₄ alkyl, C₁-C₄ haloalkyl,
 20 C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, and
 C₁-C₄ haloalkyl-S-;

R² is H, methyl, or ethyl;

25 R⁵ is OR¹⁴;

C₁-C₆ alkyl substituted with 0-3 R^{5b};

C₁-C₆ alkoxy substituted with 0-3 R^{5b};

C₂-C₆ alkenyl substituted with 0-3 R^{5b}; or

C₂-C₆ alkynyl substituted with 0-3 R^{5b};

30

R^{5a} is H or C₁-C₄ alkyl;

R^{5b}, at each occurrence, is independently selected from:

H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
NR¹⁵R¹⁶, acetyl, -S(C₁-C₄ alkyl), S(=O)(C₁-C₄
alkyl), S(=O)₂(C₁-C₄ alkyl);

C₃-C₁₀ cycloalkyl substituted with 0-3 R^{5c};

5 C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{5c};


10

R^{5c}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

15

R⁶ is H;

Ring B is selected from:



20

R¹¹, at each occurrence, is independently selected from
H, C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹,
C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹,
CF₃;

25

C₁-C₆ alkyl substituted with 0-1 R^{11a};
C₆-C₁₀ aryl substituted with 0-3 R^{11b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{11b};

R^{11a}, at each occurrence, is independently selected from H,
5 C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
CF₃, phenyl substituted with 0-3 R^{11b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
10 sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{11b};

R^{11b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
15 S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

W is -(CR⁸R^{8a})_p-;

20 p is 0, 1, or 2;

R⁸ and R^{8a}, at each occurrence, are independently selected
from H, F, methyl, and ethyl;

25 X is a bond;

phenyl substituted with 0-3 R^{Xb};
C₃-C₆ cycloalkyl substituted with 0-3 R^{Xb}; or
5 to 6 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
30 sulphur, wherein said 5 to 6 membered heterocycle
is substituted with 0-2 R^{Xb};

R^{Xb}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,

S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

Y is a bond or -(CR⁹R^{9a})_t-V-(CR⁹R^{9a})_u-;

5

t is 0, 1, or 2;

u is 0, 1, or 2;

10 R⁹ and R^{9a}, at each occurrence, are independently selected from H, F, methyl, and ethyl;

V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-,
-N(R¹⁹)-, -C(=O)NH-, or -NHC(=O)-;

15

Z is H;

C₁-C₆ alkyl substituted with 0-2 R¹²;

C₂-C₆ alkenyl substituted with 0-2 R¹²;

C₂-C₆ alkynyl substituted with 0-2 R¹²;

20 phenyl substituted with 0-4 R^{12b};

C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered

25 heterocycle is substituted with 0-3 R^{12b};

R¹² is phenyl substituted with 0-4 R^{12b};

C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or

30 5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b};

R^{12b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

5

R¹³, at each occurrence, is independently selected from H, OH, C₁-C₄ alkyl, C₁-C₄ alkoxy, Cl, F, Br, CN, NO₂, and CF₃;

10 R¹⁴, at each occurrence, is independently selected from H, phenyl, benzyl, C₁-C₆ alkyl, or C₂-C₆ alkoxyalkyl;

R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

15

R¹⁶, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

20

alternatively, -NR¹⁵R¹⁶ may be a heterocyclic ring selected from the group piperidinyl, morpholinyl, thiomorpholinyl, pyrrolidinyl, homopiperidinyl, piperazinyl, and N-methylpiperizinyl;

25

R¹⁷ is H, aryl, aryl-CH₂-, C₁-C₆ alkyl, or C₂-C₆ alkoxyalkyl;

R¹⁸, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);


30

R¹⁹, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl); and

5 R²⁰ is H or C₁-C₄;

provided that when Q is phenyl, 3-F-phenyl, benzyl, propyl, or octyl; R² is H; R⁵ is methyl, and -WXYZ is methyl, then Ring B is not


10



[5] In another alternative embodiment of the present invention of of Formula (I),

15

Ring B is



Q is C₁-C₆ alkyl substituted with 0-2 R^{1a};

20

C₆-C₁₀ aryl substituted with 0-3 R^{1b}; or

5 to 10 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{1b};

25

R^{1a}, at each occurrence, is independently selected from H, Cl, F, Br, NR¹⁹R²⁰, CF₃, C₁-C₄ alkyl, C₁-C₄ haloalkyl, and phenyl substituted with 0-3 R^{1b};

- R^{1b} , at each occurrence, is independently selected from H, OR^{14} , Cl, F, Br, NO_2 , $NR^{19}R^{20}$, CF_3 , OCF_3 , C_1-C_4 alkoxy, C_1-C_4 haloalkoxy; $-C(=O)-R^{1d}$, $-O-R^{1d}$, $-S(=O)_2-R^{1d}$, $-N(R^{19})-R^{1d}$, $-C(=O)NR^{19b}R^{1d}$, $-NR^{19b}C(=O)-R^{1d}$, $-NR^{19b}S(=O)_2-R^{1d}$, $-S(=O)_2NR^{19b}-R^{1d}$, $-C(=O)O-R^{1d}$, $-OC(=O)-R^{1d}$; C_1-C_6 alkyl substituted with 0-3 R^{1c} ; C_6-C_{10} aryl substituted with 0-3 R^{1f} ; and 5 to 14 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 14 membered heterocycle is substituted with 0-3 R^{1f} ;
- R^{1c} , at each occurrence, is independently selected from H, $-C(=O)-R^{1d}$, $-O-R^{1d}$, $-S-R^{1d}$, $-S(=O)-R^{1d}$, $-S(=O)_2-R^{1d}$, $-N(R^{19})-R^{1d}$, $-C(=O)NR^{19b}R^{1d}$, $-NR^{19b}C(=O)-R^{1d}$, $-NR^{19b}S(=O)_2-R^{1d}$, $-S(=O)_2NR^{19b}-R^{1d}$, $-NR^{19b}S(=O)-R^{1d}$, $-S(=O)NR^{19b}-R^{1d}$, $-C(=O)O-R^{1d}$, and $-OC(=O)-R^{1d}$;
- R^{1d} , at each occurrence, is independently selected from H, C_1-C_6 alkyl, and C_6-C_{10} aryl substituted with 0-3 R^{1f} ;
- R^{1f} , at each occurrence, is independently selected from H, Cl, F, Br, NO_2 , CF_3 , C_1-C_4 alkyl, C_1-C_4 haloalkyl, C_1-C_4 alkoxy, C_1-C_4 haloalkoxy, and C_1-C_4 haloalkyl-S-;
- R^5 is OR^{14} ; C_1-C_6 alkyl substituted with 0-1 R^{5b} ; C_2-C_6 alkenyl substituted with 0-1 R^{5b} ; or

C₂-C₆ alkynyl substituted with 0-1 R^{5b};

R^{5a} is H or C₁-C₄ alkyl;

5 R^{5b}, at each occurrence, is independently selected from:

H, methyl, ethyl, propyl, butyl, CF₃, OR¹⁴, Cl, F, Br,
I, =O, CN, NO₂, NR¹⁵R¹⁶; acetyl, -S(C₁-C₄ alkyl),
-S(=O)(C₁-C₄ alkyl), -S(=O)₂(C₁-C₄ alkyl),

C₃-C₇ cycloalkyl substituted with 0-3 R^{5c};

10 phenyl substituted with 0-3 R^{5c}; and

5 to 7 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle
is substituted with 0-3 R^{5c};

15

R^{5c}, at each occurrence, is independently selected from

H, OH, Cl, F, -NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)₂CH₃,
methyl, ethyl, propyl, butyl, methoxy, ethoxy,
propoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

20

R¹¹, at each occurrence, is independently selected from

H, -NR¹⁸R¹⁹, -C(=O)R¹⁷, -C(=O)OR¹⁷, -C(=O)NR¹⁸R¹⁹,
-S(=O)₂NR¹⁸R¹⁹, -CF₃;

C₁-C₆ alkyl substituted with 0-1 R^{11a};

25

phenyl substituted with 0-3 R^{11b};

C₃-C₆ carbocycle substituted with 0-3 R^{11b}; or

5 to 7 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle
is substituted with 0-3 R^{11b};

30

- R^{11a}, at each occurrence, is independently selected from H, C₁-C₄ alkyl, OR¹⁴, Cl, F, Br, CN, NO₂, NR¹⁵R¹⁶, CF₃, phenyl substituted with 0-3 R^{11b}; C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
- 5 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{11b};
- 10 R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;
- 15 W is -(CR⁸R^{8a})_p-;
- p is 0, 1, or 2;
- R⁸ and R^{8a}, at each occurrence, are independently selected
- 20 from H, F, methyl, and ethyl;
- X is a bond;
- phenyl substituted with 0-3 R^{Xb};
- C₃-C₆ cycloalkyl substituted with 0-3 R^{Xb}; or
- 25 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-2 R^{Xb};
- 30 R^{Xb}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

Y is a bond or $-(CR^9R^{9a})_t-V-(CR^9R^{9a})_u-$;

t is 0, 1, or 2;

5 u is 0, 1, or 2;

R^9 and R^{9a} , at each occurrence, are independently selected from H, F, methyl, and ethyl;

10 V is a bond, $-C(=O)-$, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$,
 $-N(R^{19})-$, $-C(=O)NH-$, or $-NHC(=O)-$;

Z is H;

C₁-C₆ alkyl substituted with 0-2 R^{12} ;

15 C₂-C₆ alkenyl substituted with 0-2 R^{12} ;

C₂-C₆ alkynyl substituted with 0-2 R^{12} ;

phenyl substituted with 0-4 R^{12b} ;

C₃-C₆ carbocycle substituted with 0-4 R^{12b} ; or

5 to 10 membered heterocycle containing 1 to 4
20 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b} ;

R^{12} is phenyl substituted with 0-4 R^{12b} ;

25 C₃-C₆ carbocycle substituted with 0-4 R^{12b} ; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b} ;

30

R^{12b} , at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R¹³, at each occurrence, is independently selected from H, OH, C₁-C₃ alkyl, C₁-C₃ alkoxy, Cl, F, Br, CN, NO₂, and CF₃;

5

R¹⁴, at each occurrence, is independently selected from H, phenyl, benzyl, C₁-C₆ alkyl, and C₂-C₆ alkoxyalkyl;

10 R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

15 R¹⁶, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

alternatively, -NR¹⁵R¹⁶ may be a heterocyclic ring selected from the group piperidinyl, morpholinyl, thiomorpholinyl, pyrrolidinyl, homopiperidinyl, 20 piperazinyl, and N-methylpiperizinyl;

R¹⁷ is H, phenyl, benzyl, C₁-C₄ alkyl, or C₂-C₄ alkoxyalkyl;


25 R¹⁸, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

30 R¹⁹, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and

R²⁰ is H, methyl, ethyl, propyl, or butyl.

[6] In another alternative embodiment of the present invention of of Formula (I),

Ring B is



Q is C₁-C₆ alkyl substituted with 0-1 R^{1a};

phenyl substituted with 0-3 R^{1b};

naphthyl substituted with 0-3 R^{1b}; or

10 5 to 10 membered heterocycle containing 1 to 2
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{1b};

15 R^{1a}, at each occurrence, is independently selected from
H, Cl, F, Br, -NR¹⁹R²⁰, -CF₃; and
phenyl substituted with 0-3 R^{1b};

20 R^{1b}, at each occurrence, is independently selected from
H, methyl, ethyl, OR¹⁴, Cl, F, Br, NO₂, NR¹⁹R²⁰, CF₃,
OCF₃, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;
-C(=O)-R^{1d}, -O-R^{1d}, -S(=O)₂-R^{1d}, -N(R¹⁹)-R^{1d},
-C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d}, -NR^{19b}S(=O)₂-R^{1d},
-S(=O)₂NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, -OC(=O)-R^{1d};
25 C₁-C₆ alkyl substituted with 0-3 R^{1c};
phenyl substituted with 0-3 R^{1f}; and
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
30 heterocycle is substituted with 0-3 R^{1f};

R^{1c} , at each occurrence, is independently selected from
H, $-C(=O)-R^{1d}$, $-O-R^{1d}$, $-S-R^{1d}$, $-S(=O)-R^{1d}$,
 $-S(=O)_2-R^{1d}$, $-N(R^{19})-R^{1d}$, $-C(=O)NR^{19b}R^{1d}$,
5 $-NR^{19b}C(=O)-R^{1d}$, $-NR^{19b}S(=O)_2-R^{1d}$, and
 $-S(=O)_2NR^{19b}-R^{1d}$;

R^{1d} , at each occurrence, is independently selected from
H, C_1-C_6 alkyl; and phenyl substituted with 0-3 R^{1f} ;
10

R^{1f} , at each occurrence, is independently selected from
H, Cl, F, Br, NO_2 , CF_3 , methyl, ethyl, propyl,
methoxy, ethoxy, propoxy, C_1-C_2 haloalkyl, and
 C_1-C_2 haloalkoxy;
15

R^5 is OR^{14} or C_1-C_4 alkyl substituted with 0-1 R^{5b} ;

R^{5a} is H;

20 R^{5b} , at each occurrence, is independently selected from:
H, methyl, ethyl, propyl, butyl, $-NR^{15}R^{16}$;
 C_3-C_7 cycloalkyl substituted with 0-3 R^{5c} ;
phenyl substituted with 0-3 R^{5c} ; or
5 to 7 membered heterocycle containing 1 to 4
25 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle
is substituted with 0-3 R^{5c} ; wherein said 5 to 7
membered heterocycle is selected from pyridinyl,
pyrimidinyl, triazinyl, furanyl, thienyl,
30 thiazolyl, pyrrolyl, piperazinyl, piperidinyl,
homopiperidinyl, pyrazolyl, imidazolyl, oxazolyl,
isoxazolyl, and tetrazolyl;

R^{5c}, at each occurrence, is independently selected from H, OH, Cl, F, -NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)₂CH₃, methyl, and methoxy;

- 5 R¹¹, at each occurrence, is independently selected from
H, NR¹⁸R¹⁹, CF₃;
C₁-C₆ alkyl substituted with 0-1 R^{11a};
phenyl substituted with 0-3 R^{11b};
C₃-C₆ carbocycle substituted with 0-3 R^{11b}; or
10 5 to 7 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle
is substituted with 0-3 R^{11b}; wherein said 5 to 7
membered heterocycle is selected from pyridinyl,
15 pyrimidinyl, triazinyl, furanyl, thienyl,
thiazolyl, pyrrolyl, piperazinyl, piperidinyl,
homopiperidinyl, pyrazolyl, imidazolyl, oxazolyl,
isoxazolyl, and tetrazolyl;
- 20 R^{11a}, at each occurrence, is independently selected from H,
methyl, ethyl, propyl, butyl, methoxy, ethoxy,
propoxy, Cl, F, Br, CN, NO₂, NR¹⁵R¹⁶, CF₃,
phenyl substituted with 0-3 R^{11b};
C₃-C₆ carbocycle substituted with 0-3 R^{11b}; or
25 5 to 7 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle
is substituted with 0-3 R^{11b}; wherein said 5 to 7
membered heterocycle is selected from pyridinyl,
30 pyrimidinyl, triazinyl, furanyl, thienyl,
thiazolyl, pyrrolyl, piperazinyl, piperidinyl,
homopiperidinyl, pyrazolyl, imidazolyl, oxazolyl,
isoxazolyl, and tetrazolyl;

R^{11b}, at each occurrence, is independently selected from H,
OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃,
S(=O)₂CH₃, methyl, ethyl, propyl, butyl, methoxy,
ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂
5 haloalkoxy;

W is -(CHR⁸)_p-;

p is 0 or 1;

R⁸ is H, methyl, or ethyl;

X is a bond;

phenyl substituted with 0-2 R^{Xb};

15 C₃-C₆ cycloalkyl substituted with 0-3 R^{Xb}; or
5 to 6 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 6 membered heterocycle
is substituted with 0-1 R^{Xb};

20 R^{Xb}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

25 Y is a bond, -V-, -CH₂-V-, -V-CH₂-, or -CH₂-V-CH₂-;

V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-,
-N(R¹⁹)-, -C(=O)NH-, or -NHC(=O)-;

30 Z is H;

C₁-C₆ alkyl substituted with 0-2 R¹²;

C₂-C₆ alkenyl substituted with 0-2 R¹²;

C₂-C₆ alkynyl substituted with 0-2 R¹²;

- phenyl substituted with 0-4 R^{12b};
C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
5 sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b};
- R¹² is phenyl substituted with 0-4 R^{12b};
C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or
10 5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b};
- 15 R^{12b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;
- 20 R¹³, at each occurrence, is independently selected from
H, methyl, ethyl, methoxy, ethoxy, Cl, F, Br, NO₂,
or CF₃;
- R¹⁴, at each occurrence, is independently selected from H,
25 phenyl, benzyl, C₁-C₄ alkyl, or C₂-C₄ alkoxyalkyl;
- R¹⁵, at each occurrence, is independently selected from H,
methyl, ethyl, propyl, butyl, phenyl, benzyl, and
phenethyl;
- 30 R¹⁶, at each occurrence, is independently selected from H,
OH, methyl, ethyl, propyl, butyl, phenyl, benzyl,
phenethyl, CH₃CH₂C(=O)-, CH₃C(=O)-, CH₃CH₂OC(=O)-,
CH₃OC(=O)-, CH₃CH₂S(=O)₂- and CH₃S(=O)₂-;
- 35

R¹⁸, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl;


5 R¹⁹, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, and butyl; and

R²⁰ is H, methyl, ethyl, propyl, or butyl.

10

[7] In another alternative embodiment of the present invention of of Formula (I),

Ring B is



Q is C₁-C₄ alkyl substituted with 0-1 R^{1a};

phenyl substituted with 0-3 R^{1b};

naphthyl substituted with 0-3 R^{1b}; or

20

5 to 10 membered heterocycle containing 1 to 2

heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered

heterocycle is substituted with 0-3 R^{1b}; wherein

said 5 to 10 membered heterocycle is selected

25

from pyridinyl, quinolinyl, pyrimidinyl,

triazinyl, furanyl, thienyl, thiazolyl,

imidazolyl, oxazolyl, and isoxazolyl;

R^{1a}, at each occurrence, is independently selected from

30

H, Cl, F, Br, CF₃, and

phenyl substituted with 0-3 R^{1b};

R^{1b}, at each occurrence, is independently selected from
H, methyl, ethyl, OR¹⁴, Cl, F, Br, NO₂, NR¹⁹R²⁰, CF₃,
OCF₃, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy; -C(=O)-R^{1d},
-O-R^{1d}, -S(=O)₂-R^{1d}, -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d},
5 -NR^{19b}C(=O)-R^{1d}, -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d},
-C(=O)O-R^{1d}, -OC(=O)-R^{1d},
C₁-C₆ alkyl substituted with 0-3 R^{1c};
phenyl substituted with 0-3 R^{1f}; and
5 to 6 membered heterocycle containing 1 to 4
10 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 6 membered heterocycle
is substituted with 0-3 R^{1f};

R^{1c}, at each occurrence, is independently selected from
15 H, -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d}, -S(=O)-R^{1d},
-S(=O)₂-R^{1d}, -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d},
-NR^{19b}C(=O)-R^{1d}, -NR^{19b}S(=O)₂-R^{1d}, and
-S(=O)₂NR^{19b}-R^{1d},

20 R^{1d}, at each occurrence, is independently selected from
H, C₁-C₆ alkyl; and phenyl substituted with 0-3 R^{1f};

R^{1f}, at each occurrence, is independently selected from
H, Cl, F, Br, NO₂, CF₃, -OCF₃, methyl, ethyl, methoxy,
25 and ethoxy;

R⁵ is methyl substituted with 0-1 R^{5b};
ethyl substituted with 0-1 R^{5b};
propyl substituted with 0-1 R^{5b}; or
30 butyl substituted with 0-1 R^{5b};

R^{5a} is H;

R^{5b}, at each occurrence, is independently selected from:

H, methyl, ethyl, and NR¹⁵R¹⁶;

R¹¹, at each occurrence, is independently selected from

5 H, NR¹⁸R¹⁹, CF₃;

C₁-C₄ alkyl substituted with 0-1 R^{11a};

phenyl substituted with 0-3 R^{11b};

C₃-C₆ carbocycle substituted with 0-3 R^{11b}; or

5 to 7 membered heterocycle containing 1 to 4
10 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle
is substituted with 0-3 R^{11b}; wherein said 5 to 7
membered heterocycle is selected from pyridinyl,
pyrimidinyl, triazinyl, furanyl, thienyl,
15 thiazolyl, pyrrolyl, piperazinyl, piperidinyl,
homopiperidinyl, pyrazolyl, imidazolyl, oxazolyl,
isoxazolyl, and tetrazolyl;

R^{11a}, at each occurrence, is independently selected from H,

20 methyl, ethyl, propyl, butyl, methoxy, ethoxy,

propoxy, Cl, F, NR¹⁵R¹⁶, CF₃;

phenyl substituted with 0-3 R^{11b};

C₃-C₆ carbocycle substituted with 0-3 R^{11b}; or

5 to 7 membered heterocycle containing 1 to 4
25 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle
is substituted with 0-3 R^{11b}; wherein said 5 to 7
membered heterocycle is selected from pyridinyl,
pyrimidinyl, triazinyl, furanyl, thienyl,
30 thiazolyl, pyrrolyl, piperazinyl, piperidinyl,
homopiperidinyl, pyrazolyl, imidazolyl, oxazolyl,
isoxazolyl, and tetrazolyl;

R^{11b}, at each occurrence, is independently selected from

H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, methoxy, and ethoxy;

W is a bond, -CH₂-, or -CH(CH₃)-;

5

X is a bond;

phenyl substituted with 0-1 R^{Xb};

C₃-C₆ cycloalkyl substituted with 0-3 R^{Xb}; or

5 to 6 membered heterocycle containing 1 to 3

10 heteroatoms selected from nitrogen, oxygen, and sulphur; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, 15 oxazolyl, and isoxazolyl;

Y is a bond, -V-, -CH₂-V-, or -V-CH₂-;

V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-,

20 -N(R¹⁹)-, -C(=O)NH-, or -NHC(=O)-;

Z is H;

C₁-C₆ alkyl substituted with 0-2 R¹²;

C₂-C₆ alkenyl substituted with 0-2 R¹²;

25 C₂-C₆ alkynyl substituted with 0-2 R¹²;

phenyl substituted with 0-4 R^{12b};

C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4

30 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

R¹² is phenyl substituted with 0-4 R^{12b};

C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b};

5

R^{12b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, NR¹⁵R¹⁶, CF₃, OCF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, methyl, ethyl, propyl, butyl,
methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂
haloalkoxy;

10

R¹³, at each occurrence, is independently selected from
H, methyl, methoxy, Cl, F, Br, and CF₃;

15 R¹⁴, at each occurrence, is independently selected from H,
phenyl, benzyl, methyl, ethyl, propyl, and butyl;

R¹⁵, at each occurrence, is independently selected from H,
methyl, ethyl, propyl, or butyl;

20

R¹⁶, at each occurrence, is independently selected from H,
OH, methyl, ethyl, propyl, butyl, phenyl, benzyl, and
phenethyl;

25 R¹⁸, at each occurrence, is independently selected from H,
methyl, ethyl, propyl, butyl, phenyl, benzyl, and
phenethyl;

30 R¹⁹, at each occurrence, is independently selected from H,
OH, methyl, ethyl, propyl, and butyl; and

R²⁰ is H, methyl, ethyl, propyl, or butyl.

[8] In another alternative embodiment of the present
35 invention of of Formula (I),

Q is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH₂CH₂CH₂CH₃, -CH(CH₃)₂,
-CH(CH₃)CH₂CH₃, -CH₂CH(CH₃)₂,

-CH₂CF₃, -CH₂CH₂CF₃, -CH₂CH₂CH₂CF₃,

5

phenyl-, 4-tBu-phenyl-, 4-iPr-phenyl-, 4-Et-phenyl-,

2-F-phenyl-, 3-F-phenyl-, 4-F-phenyl-,

2-Cl-phenyl-, 3-Cl-phenyl-, 4-Cl-phenyl-,

2-Br-phenyl-, 3-Br-phenyl-, 4-Br-phenyl-,

10 2-NO₂-phenyl-, 3-NO₂-phenyl-, 4-NO₂-phenyl-,

2-CH₃-phenyl-, 3-CH₃-phenyl-, 4-CH₃-phenyl-,

2-CH₃O-phenyl-, 3-CH₃O-phenyl-, 4-CH₃O-phenyl-,

2-CF₃-phenyl-, 3-CF₃-phenyl-, 4-CF₃-phenyl-,

2-CF₃O-phenyl-, 3-CF₃O-phenyl-, 4-CF₃O-phenyl-,

15 2-CH₃CONH-phenyl, 3-CH₃CONH-phenyl, 4-CH₃CONH-phenyl,

2-methoxycarbonyl-phenyl-, 4-phenyl-phenyl-,

2,3-diF-phenyl-, 2,4-diF-phenyl-, 2,5-diF-phenyl-,

2,6-diF-phenyl-, 3,4-diF-phenyl-, 3,5-diF-phenyl-,

20 2,3-diCl-phenyl-, 2,4-diCl-phenyl-, 2,5-diCl-phenyl-,

2,6-diCl-phenyl-, 3,4-diCl-phenyl-, 3,5-diCl-phenyl-,

3-F-4-Cl-phenyl-, 3-F-5-Cl-phenyl-, 3-Cl-4-F-phenyl-,

2,3-diMe-phenyl-, 2,4-diMe-phenyl-, 2,5-diMe-phenyl-,

2,6-diMe-phenyl-, 3,4-diMe-phenyl-, 3,5-diMe-phenyl-,

25 2,3-diMeO-phenyl-, 2,4-diMeO-phenyl-, 2,5-diMeO-phenyl-,

2,6-diMeO-phenyl-, 3,4-diMeO-phenyl-, 3,5-diMeO-phenyl-,

2,3-diCF₃-phenyl-, 2,4-diCF₃-phenyl-, 2,5-diCF₃-phenyl-,

2,6-diCF₃-phenyl-, 3,4-diCF₃-phenyl-, 3,5-diCF₃-phenyl-,

2-NO₂-4-CF₃-phenyl-, 2-Me-5-Cl-phenyl-,

30 2,4,6-triMe-phenyl,

benzyl-, naphth-1-yl-, naphth-2-yl-, beta-styrene-,

furanyl-, thienyl-, pyridyl-, thiazolyl-, imidazol-1-yl-

, oxazolyl-, isoxazolyl-, quinolin-8-yl-,


35

3-methyl-isoxazol-4-yl-,

- 3,5-dimethyl-isoxazol-4-yl-,
 3-bromo-5-chloro-thiophen-2-yl-,
 2,3-dichlorothiophen-5-yl-,
 4-bromo-5-chlorothiophen-2-yl-,
 5-[(benzoylamino)methyl]-thiophen-2-yl-,
 4-phenylsulfonylthiophen-2-yl-,
 5-(phenylsulfonyl)thiophen-2-yl-,
 2-(1-methyl-(5-trifluoromethyl)pyrazole)thiophen-5-yl-,
 5-(2-pyridyl)thiophen-2-yl-,
 1-methyl-5-(trifluoromethyl)imidazol-3-yl-,
 2-(2-methylthio-pyrimidin-3-yl)-thiophen-5-yl-, or
 dibenzofuran-2-yl-;
- R⁵ is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃,
 -CH(CH₃)CH₂CH₃, -CH₂CH(CH₃)₂, -CH₂C(CH₃)₃,
 -CH₂CH₂CH₂CH₂CH₃, -CH(CH₃)CH₂CH₂CH₃, -CH₂CH(CH₃)CH₂CH₃,
 -CH₂CH₂CH(CH₃)₂, -CH(CH₂CH₃)₂,
- cyclopropyl-CH₂-, cyclobutyl-CH₂-, cyclopentyl-CH₂-,
 cyclohexyl-CH₂-, (2-CH₃-cyclopropyl)CH₂-,
 (3-CH₃-cyclobutyl)CH₂-, cyclopropyl-CH₂CH₂-,
 cyclobutyl-CH₂CH₂-, cyclopentyl-CH₂CH₂-,
 cyclohexyl-CH₂CH₂-, (2-CH₃-cyclopropyl)CH₂CH₂-,
 (3-CH₃-cyclobutyl)CH₂CH₂-,
- phenyl-CH₂-, (2-F-phenyl)CH₂-, (3-F-phenyl)CH₂-,
 (4-F-phenyl)CH₂-, furanyl-CH₂-, thienyl-CH₂-,
 pyridyl-CH₂-, 1-imidazolyl-CH₂-, oxazolyl-CH₂-,
 isoxazolyl-CH₂-,
- phenyl-CH₂CH₂-, (2-F-phenyl)CH₂CH₂-, (3-F-phenyl)CH₂CH₂-,
 (4-F-phenyl)CH₂CH₂-, furanyl-CH₂CH₂-, thienyl-CH₂CH₂-,
 pyridyl-CH₂CH₂-, 1-imidazolyl-CH₂CH₂-, oxazolyl-CH₂CH₂-,
 or isoxazolyl-CH₂CH₂-,

W is a bond, $-\text{CH}_2-$, or $-\text{CH}(\text{CH}_3)-$;

X is a bond;



Y is a bond, $-\text{CH}_2-\text{V}-$, $-\text{V}-$, or $-\text{V}-\text{CH}_2-$;

10

V is a bond, $-\text{C}(=\text{O})-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(=\text{O})-$, $-\text{S}(=\text{O})_2-$, $-\text{NH}-$,
 $-\text{N}(\text{CH}_3)-$, $-\text{C}(=\text{O})\text{NH}-$, or $-\text{NHC}(=\text{O})-$,

Z is H, $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)_2$,

15

$-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{C}(\text{CH}_3)_3$,
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$,
 $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_2\text{CH}_3)_2$, $-\text{CH}_2\text{C}(\text{CH}_3)_3$, $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_3$,

$-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$,

20

$-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$,
 $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)_2$,

$-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_3$, $-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$,

$-\text{CF}_3$, $-\text{CH}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CF}_3$,

25

cyclopropyl-, (cyclopropyl) CH_2- , (cyclopropyl) CH_2CH_2- ,

cyclobutyl-, (cyclobutyl) CH_2- , (cyclobutyl) CH_2CH_2- ,

cyclopentyl-, (cyclopentyl) CH_2- , (cyclopentyl) CH_2CH_2- ,

cyclohexyl-, (cyclohexyl) CH_2- , (cyclohexyl) CH_2CH_2- ,

30

phenyl, 2-F-phenyl, 3-F-phenyl, 4-F-phenyl, 2-Cl-phenyl,
 3-Cl-phenyl, 4-Cl-phenyl, 2,3-diF-phenyl,
 2,4-diF-phenyl, 2,5-diF-phenyl, 2,6-diF-phenyl,

3,4-diF-phenyl, 3,5-diF-phenyl, 2,3-diCl-phenyl,
2,4-diCl-phenyl, 2,5-diCl-phenyl, 2,6-diCl-phenyl,
3,4-diCl-phenyl, 3,5-diCl-phenyl, 3-F-4-Cl-phenyl,
3-F-5-Cl-phenyl, 3-Cl-4-F-phenyl, 2-MeO-phenyl,
5 3-MeO-phenyl, 4-MeO-phenyl, 2-Me-phenyl, 3-Me-phenyl,
4-Me-phenyl, 2-MeS-phenyl, 3-MeS-phenyl, 4-MeS-phenyl,
2-CF₃O-phenyl, 3-CF₃O-phenyl, 4-CF₃O-phenyl, furanyl,
thienyl, pyridyl, 2-Me-pyridyl, 3-Me-pyridyl,
4-Me-pyridyl, 1-imidazolyl, oxazolyl, isoxazolyl,
10 1-benzimidazolyl, morpholino, N-piperidiny,
phenyl-CH₂-, (2-F-phenyl)CH₂-, (3-F-phenyl)CH₂-,
(4-F-phenyl)CH₂-, (2-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂-,
(4-Cl-phenyl)CH₂-, (2,3-diF-phenyl)CH₂-,
(2,4-diF-phenyl)CH₂-, (2,5-diF-phenyl)CH₂-,
15 (2,6-diF-phenyl)CH₂-, (3,4-diF-phenyl)CH₂-,
(3,5-diF-phenyl)CH₂-, (2,3-diCl-phenyl)CH₂-,
(2,4-diCl-phenyl)CH₂-, (2,5-diCl-phenyl)CH₂-,
(2,6-diCl-phenyl)CH₂-, (3,4-diCl-phenyl)CH₂-,
(3,5-diCl-phenyl)CH₂-, (3-F-4-Cl-phenyl)CH₂-,
20 (3-F-5-Cl-phenyl)CH₂-, (3-Cl-4-F-phenyl)CH₂-,
(2-MeO-phenyl)CH₂-, (3-MeO-phenyl)CH₂-,
(4-MeO-phenyl)CH₂-, (2-Me-phenyl)CH₂-,
(3-Me-phenyl)CH₂-, (4-Me-phenyl)CH₂-, (2-MeS-phenyl)CH₂-,
, (3-MeS-phenyl)CH₂-, 4-MeS-phenyl)CH₂-,
25 (2-CF₃O-phenyl)CH₂-, (3-CF₃O-phenyl)CH₂-,
(4-CF₃O-phenyl)CH₂-, (furanyl)CH₂-, (thienyl)CH₂-,
(pyridyl)CH₂-, (2-Me-pyridyl)CH₂-, (3-Me-pyridyl)CH₂-,
(4-Me-pyridyl)CH₂-, (1-imidazolyl)CH₂-, (oxazolyl)CH₂-,
(isoxazolyl)CH₂-, (1-benzimidazolyl)CH₂-,
30 morpholino)CH₂-, (N-piperidiny)CH₂-,
phenyl-CH₂CH₂-, (phenyl)₂CHCH₂-, (2-F-phenyl)CH₂CH₂-,
(3-F-phenyl)CH₂CH₂-, (4-F-phenyl)CH₂CH₂-,
(2-Cl-phenyl)CH₂CH₂-, (3-Cl-phenyl)CH₂CH₂-,
(4-Cl-phenyl)CH₂CH₂-, (2,3-diF-phenyl)CH₂CH₂-,
35 (2,4-diF-phenyl)CH₂CH₂-, (2,5-diF-phenyl)CH₂CH₂-,

(2,6-diF-phenyl)CH₂CH₂-, (3,4-diF-phenyl)CH₂CH₂-,
 (3,5-diF-phenyl)CH₂CH₂-, (2,3-diCl-phenyl)CH₂CH₂-,
 (2,4-diCl-phenyl)CH₂CH₂-, (2,5-diCl-phenyl)CH₂CH₂-,
 (2,6-diCl-phenyl)CH₂CH₂-, (3,4-diCl-phenyl)CH₂CH₂-,
 5 (3,5-diCl-phenyl)CH₂CH₂-, (3-F-4-Cl-phenyl)CH₂CH₂-,
 (3-F-5-Cl-phenyl)CH₂CH₂-, (3-Cl-4-F-phenyl)CH₂CH₂-,
 (2-MeO-phenyl)CH₂CH₂-, (3-MeO-phenyl)CH₂CH₂-,
 (4-MeO-phenyl)CH₂CH₂-, (2-Me-phenyl)CH₂CH₂-,
 (3-Me-phenyl)CH₂CH₂-, (4-Me-phenyl)CH₂CH₂-,
 10 (2-MeS-phenyl)CH₂CH₂-, (3-MeS-phenyl)CH₂CH₂-,
 (4-MeS-phenyl)CH₂CH₂-, (2-CF₃O-phenyl)CH₂CH₂-,
 (3-CF₃O-phenyl)CH₂CH₂-, (4-CF₃O-phenyl)CH₂CH₂-,
 (furanyl)CH₂CH₂-, (thienyl)CH₂CH₂-, (pyridyl)CH₂CH₂-,
 (2-Me-pyridyl)CH₂CH₂-, (3-Me-pyridyl)CH₂CH₂-,
 15 (4-Me-pyridyl)CH₂CH₂-, (imidazolyl)CH₂CH₂-,
 (oxazolyl)CH₂CH₂-, (isoxazolyl)CH₂CH₂-,
 (benzimidazolyl)CH₂CH₂-, (cyclopropyl)CH₂CH₂-,
 (cyclobutyl)CH₂CH₂-, (cyclopentyl)CH₂CH₂-,
 (cyclohexyl)CH₂CH₂-, (morpholino)CH₂CH₂-, or
 20 (N-piperidinyl)CH₂CH₂-;

R¹¹, at each occurrence, is independently selected from H,
 methyl, ethyl, propyl, butyl, phenyl, benzyl, phenethyl,
 25 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,
 cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl,
 cyclohexylmethyl, cyclopropylethyl, cyclobutylethyl,
 cyclopentylethyl, cyclohexylethyl, 4-F-phenyl,
 4-Cl-phenyl, 4-CH₃-phenyl, 4-CF₃-phenyl, 4-CH₃O-phenyl,
 30 4-CF₃O-phenyl, 3-F-phenyl, 3-Cl-phenyl, 3-CH₃-phenyl,
 3-CF₃-phenyl, 3-CH₃O-phenyl, 3-CF₃O-phenyl, 2-F-phenyl,
 2-Cl-phenyl, 2-CH₃-phenyl, 2-CF₃-phenyl, 2-CH₃O-phenyl,
 2-CF₃O-phenyl, (4-F-phenyl)methyl-, (4-Cl-phenyl)methyl-,
 ,
 35 (4-CH₃-phenyl)methyl-, (4-CF₃-phenyl)methyl-,

(4-CH₃O-phenyl)methyl-, (4-CF₃O-phenyl)methyl-,
 (3-F-phenyl)methyl-, (3-Cl-phenyl)methyl-,
 (3-CH₃-phenyl)methyl-, (3-CF₃-phenyl)methyl-,
 (3-CH₃O-phenyl)methyl-, (3-CF₃O-phenyl)methyl-,
 5 (2-F-phenyl)methyl-, (2-Cl-phenyl)methyl-,
 (2-CH₃-phenyl)methyl-, (2-CF₃-phenyl)methyl-,
 (2-CH₃O-phenyl)methyl-, (2-CF₃O-phenyl)methyl-,
 2-pyridyl-, 3-pyridyl-, 4-pyridyl-,
 1-piperidinyl, 1-homopiperidinyl, and 1-morpholino; and

10

R¹³, at each occurrence, is independently selected from H,
 F, Cl, and methoxy.


[9] In another alternative embodiment of the present
 15 invention of of Formula (I),


Q is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, phenyl-,
 2-F-phenyl-, 2-Cl-phenyl-, 2-Br-phenyl-,
 2-NO₂-phenyl-, 2-CH₃-phenyl-, 2-CH₃CH₂-phenyl-,
 20 2-CH₃O-phenyl-, 2-CF₃-phenyl-, 2-CF₃O-phenyl-,
 2-CH₃CONH-phenyl, or 3,5-dimethyl-isoxazol-4-yl-;

R⁵ is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃,
 -CH(CH₃)CH₂CH₃, -CH₂CH(CH₃)₂, -CH₂C(CH₃)₃,
 25 -CH₂CH₂CH₂CH₂CH₃, -CH(CH₃)CH₂CH₂CH₃, -CH₂CH(CH₃)CH₂CH₃,
 -CH₂CH₂CH(CH₃)₂, -CH(CH₂CH₃)₂,

W is a bond, -CH₂-, or -CH(CH₃)-;

30 X is a bond;





Y is a bond, $-\text{CH}_2\text{-V-}$, $-\text{V-}$, or $-\text{V-CH}_2\text{-}$;

5 V is a bond, $-\text{C}(=\text{O})\text{-}$, $-\text{O-}$, $-\text{S-}$, $-\text{S}(=\text{O})\text{-}$, $-\text{S}(=\text{O})_2\text{-}$, $-\text{NH-}$, $-\text{N}(\text{CH}_3)\text{-}$, $-\text{C}(=\text{O})\text{NH-}$, or $-\text{NHC}(=\text{O})\text{-}$,

Z is H, $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)_2$,
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{C}(\text{CH}_3)_3$,
 10 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$,
 $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_2\text{CH}_3)_2$, $-\text{CH}_2\text{C}(\text{CH}_3)_3$, $-\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_3$,
 $-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$,
 $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$,
 15 $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}(\text{CH}_2\text{CH}_3)_2$,
 $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_3$, $-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$,
 $-\text{CF}_3$, $-\text{CH}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CF}_3$,
 20 cyclopropyl-, (cyclopropyl) $\text{CH}_2\text{-}$, (cyclopropyl) $\text{CH}_2\text{CH}_2\text{-}$,
 cyclobutyl-, (cyclobutyl) $\text{CH}_2\text{-}$, (cyclobutyl) $\text{CH}_2\text{CH}_2\text{-}$,
 cyclopentyl-, (cyclopentyl) $\text{CH}_2\text{-}$, (cyclopentyl) $\text{CH}_2\text{CH}_2\text{-}$,
 cyclohexyl-, (cyclohexyl) $\text{CH}_2\text{-}$, (cyclohexyl) $\text{CH}_2\text{CH}_2\text{-}$,
 phenyl, 2-F-phenyl, 3-F-phenyl, 4-F-phenyl, 2-Cl-phenyl,
 25 3-Cl-phenyl, 4-Cl-phenyl, 2,3-diF-phenyl,
 2,4-diF-phenyl, 2,5-diF-phenyl, 2,6-diF-phenyl,
 3,4-diF-phenyl, 3,5-diF-phenyl, 2,3-diCl-phenyl,
 2,4-diCl-phenyl, 2,5-diCl-phenyl, 2,6-diCl-phenyl,
 3,4-diCl-phenyl, 3,5-diCl-phenyl, 3-F-4-Cl-phenyl,
 30 3-F-5-Cl-phenyl, 3-Cl-4-F-phenyl, 2-MeO-phenyl,
 3-MeO-phenyl, 4-MeO-phenyl, 2-Me-phenyl, 3-Me-phenyl,
 4-Me-phenyl, 2-MeS-phenyl, 3-MeS-phenyl, 4-MeS-phenyl,
 2-CF₃O-phenyl, 3-CF₃O-phenyl, 4-CF₃O-phenyl, furanyl,
 thienyl, pyridyl, 2-Me-pyridyl, 3-Me-pyridyl,

4-Me-pyridyl, 1-imidazolyl, oxazolyl, isoxazolyl,
 1-benzimidazolyl, morpholino, N-piperidiny,
 phenyl-CH₂-, (2-F-phenyl)CH₂-, (3-F-phenyl)CH₂-,
 (4-F-phenyl)CH₂-, (2-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂-,
 5 (4-Cl-phenyl)CH₂-, (2,3-diF-phenyl)CH₂-,
 (2,4-diF-phenyl)CH₂-, (2,5-diF-phenyl)CH₂-,
 (2,6-diF-phenyl)CH₂-, (3,4-diF-phenyl)CH₂-,
 (3,5-diF-phenyl)CH₂-, (2,3-diCl-phenyl)CH₂-,
 (2,4-diCl-phenyl)CH₂-, (2,5-diCl-phenyl)CH₂-,
 10 (2,6-diCl-phenyl)CH₂-, (3,4-diCl-phenyl)CH₂-,
 (3,5-diCl-phenyl)CH₂-, (3-F-4-Cl-phenyl)CH₂-,
 (3-F-5-Cl-phenyl)CH₂-, (3-Cl-4-F-phenyl)CH₂-,
 (2-MeO-phenyl)CH₂-, (3-MeO-phenyl)CH₂-,
 (4-MeO-phenyl)CH₂-, (2-Me-phenyl)CH₂-,
 15 (3-Me-phenyl)CH₂-, (4-Me-phenyl)CH₂-, (2-MeS-phenyl)CH₂-,
 (3-MeS-phenyl)CH₂-, (4-MeS-phenyl)CH₂-,
 (2-CF₃O-phenyl)CH₂-, (3-CF₃O-phenyl)CH₂-,
 (4-CF₃O-phenyl)CH₂-, (furanyl)CH₂-, (thienyl)CH₂-,
 (pyridyl)CH₂-, (2-Me-pyridyl)CH₂-, (3-Me-pyridyl)CH₂-,
 20 (4-Me-pyridyl)CH₂-, (1-imidazolyl)CH₂-, (oxazolyl)CH₂-,
 (isoxazolyl)CH₂-, (1-benzimidazolyl)CH₂-,
 morpholino)CH₂-, (N-piperidiny)CH₂-,
 phenyl-CH₂CH₂-, (phenyl)₂CHCH₂-, (2-F-phenyl)CH₂CH₂-,
 (3-F-phenyl)CH₂CH₂-, (4-F-phenyl)CH₂CH₂-,
 25 (2-Cl-phenyl)CH₂CH₂-, (3-Cl-phenyl)CH₂CH₂-,
 (4-Cl-phenyl)CH₂CH₂-, (2,3-diF-phenyl)CH₂CH₂-,
 (2,4-diF-phenyl)CH₂CH₂-, (2,5-diF-phenyl)CH₂CH₂-,
 (2,6-diF-phenyl)CH₂CH₂-, (3,4-diF-phenyl)CH₂CH₂-,
 (3,5-diF-phenyl)CH₂CH₂-, (2,3-diCl-phenyl)CH₂CH₂-,
 30 (2,4-diCl-phenyl)CH₂CH₂-, (2,5-diCl-phenyl)CH₂CH₂-,
 (2,6-diCl-phenyl)CH₂CH₂-, (3,4-diCl-phenyl)CH₂CH₂-,
 (3,5-diCl-phenyl)CH₂CH₂-, (3-F-4-Cl-phenyl)CH₂CH₂-,
 (3-F-5-Cl-phenyl)CH₂CH₂-, (3-Cl-4-F-phenyl)CH₂CH₂-,
 (2-MeO-phenyl)CH₂CH₂-, (3-MeO-phenyl)CH₂CH₂-,


- (4-MeO-phenyl)CH₂CH₂-, (2-Me-phenyl)CH₂CH₂-,
 (3-Me-phenyl)CH₂CH₂-, (4-Me-phenyl)CH₂CH₂-,
 (2-MeS-phenyl)CH₂CH₂-, (3-MeS-phenyl)CH₂CH₂-,
 (4-MeS-phenyl)CH₂CH₂-, (2-CF₃O-phenyl)CH₂CH₂-,
 5 (3-CF₃O-phenyl)CH₂CH₂-, (4-CF₃O-phenyl)CH₂CH₂-,
 (furanyl)CH₂CH₂-, (thienyl)CH₂CH₂-, (pyridyl)CH₂CH₂-,
 (2-Me-pyridyl)CH₂CH₂-, (3-Me-pyridyl)CH₂CH₂-,
 (4-Me-pyridyl)CH₂CH₂-, (imidazolyl)CH₂CH₂-,
 (oxazolyl)CH₂CH₂-, (isoxazolyl)CH₂CH₂-,
 10 (benzimidazolyl)CH₂CH₂-, (cyclopropyl)CH₂CH₂-,
 (cyclobutyl)CH₂CH₂-, (cyclopentyl)CH₂CH₂-,
 (cyclohexyl)CH₂CH₂-, (morpholino)CH₂CH₂-, or
 (N-piperidinyl)CH₂CH₂-;
- 15 R¹¹, at each occurrence, is independently selected from H,
 methyl, ethyl, propyl, butyl, phenyl, benzyl, phenethyl,
 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,
 cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl,
 cyclohexylmethyl, cyclopropylethyl, cyclobutylethyl,
 20 cyclopentylethyl, cyclohexylethyl, 4-F-phenyl,
 4-Cl-phenyl, 4-CH₃-phenyl, 4-CF₃-phenyl, 4-CH₃O-phenyl,
 4-CF₃O-phenyl, 3-F-phenyl, 3-Cl-phenyl, 3-CH₃-phenyl,
 3-CF₃-phenyl, 3-CH₃O-phenyl, 3-CF₃O-phenyl, 2-F-phenyl,
 2-Cl-phenyl, 2-CH₃-phenyl, 2-CF₃-phenyl, 2-CH₃O-phenyl,
 25 2-CF₃O-phenyl, (4-F-phenyl)methyl-, (4-Cl-phenyl)methyl-,
 (4-CH₃-phenyl)methyl-, (4-CF₃-phenyl)methyl-,
 (4-CH₃O-phenyl)methyl-, (4-CF₃O-phenyl)methyl-,
 (3-F-phenyl)methyl-, (3-Cl-phenyl)methyl-,
 30 (3-CH₃-phenyl)methyl-, (3-CF₃-phenyl)methyl-,
 (3-CH₃O-phenyl)methyl-, (3-CF₃O-phenyl)methyl-,
 (2-F-phenyl)methyl-, (2-Cl-phenyl)methyl-,
 (2-CH₃-phenyl)methyl-, (2-CF₃-phenyl)methyl-,
 (2-CH₃O-phenyl)methyl-, (2-CF₃O-phenyl)methyl-,
 35 2-pyridyl-, 3-pyridyl-, 4-pyridyl-,

1-piperidinyl, 1-homopiperidinyl, and 1-morpholino; and

R¹³, at each occurrence, is independently selected from H, F, Cl, and methoxy.

5

[10] In another alternative embodiment, the present invention provides a novel compound of Formula (Ic):




10

(Ic)

or a pharmaceutically acceptable salt form or prodrug thereof.

15

[11] In another alternative embodiment, the present invention provides a novel compound of Formula (Id):




20

(Id)

or a pharmaceutically acceptable salt form or prodrug thereof.

25

[12] In another alternative embodiment, the present invention provides a novel compound of Formula (Ia);



(Ia)

or a pharmaceutically acceptable salt or prodrug thereof.

[13] In another alternative embodiment of the present
5 invention of Formula (I), R⁵ is OR¹⁴.

[14] In another alternative embodiment of the present
invention of Formula (I), R⁵ is C₁-C₄ alkoxy substituted
with 0-3 R^{5b}.

10

[15] In a preferred embodiment the present invention
provides compound of Formula (I), (Ia), (Ic), or (Id)
wherein Q is 3,5-dimethyl-isoxazol-4-yl-.

15 In a preferred embodiment Q is -CH₃, -CH₂CH₃,
-CH₂CH₂CH₃, -CH(CH₃)₂, phenyl-, 2-F-phenyl-, 2-Cl-phenyl-,
2-Br-phenyl-, 2-NO₂-phenyl-, 2-CH₃-phenyl-,
2-CH₃CH₂-phenyl-, 2-CH₃O-phenyl-, 2-CF₃-phenyl-,
2-CF₃O-phenyl-, 2-CH₃CONH-phenyl, or
20 3,5-dimethyl-isoxazol-4-yl-.

In a preferred embodiment Q is a substituted phenyl as
disclosed herein (preferably ortho substituted) or a 5 to 6
membered heterocyclic ring as disclosed herein (preferably
25 substituted); for example, 2-F-phenyl-, 2-Cl-phenyl-,
2-Br-phenyl-, 2-NO₂-phenyl-, 2-CH₃-phenyl-,
2-CH₃CH₂-phenyl-, 2-CH₃O-phenyl-, 2-CF₃-phenyl-,
2-CF₃O-phenyl-, 2-CH₃CONH-phenyl, or
3,5-dimethyl-isoxazol-4-yl-.

30

In a preferred embodiment Q is not a trihalomethyl
group, preferably not trifluoromethyl.

In a preferred embodiment when R⁵ and/or R^{5a} form an
35 alkyl or alkenyl moiety, optionally substituted, of five

carbons or less, then Q is not a haloalkyl, preferably not trifluoromethyl.

5

It is understood that any and all embodiments of the present invention may be taken in conjunction with any other embodiment to describe additional even more preferred embodiments of the present invention.

10

[16] In another alternative embodiment, the present invention provides a compound selected from:

15 2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

20 2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-propionamide;

25 2-(3-bromo-5-chloro-thiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

30 2-(naphthalene-1-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

35 2-(5-dimethylamino naphthalene-1-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

40 2-(naphthalene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

2-(2-acetamido-4-methylthiazole-5-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

5 2-(thiophene-2-sulfonylamino)-4-methyl-pentanoic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

10 2-(quinoline-8-sulfonylamino)-4-methyl-pentanoic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

15 2-(phenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

20 2-(2,5-dichlorophenyl sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

2 2-(mesitylene-2-sulfonylamino)-4-methyl-pentanoic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

25 2-(3-nitrophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

30 2-(4-bromophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

35 2-(4-fluorophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

2-(4-chlorophenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

5 2-(4-acetamidophenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

10 2-(4-nitrophenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

15 2-(4-methoxyphenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

20 2-(4-tert-butylphenyl-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

2-(p-toluene-sulfonylamino)-4-methyl-pentanoic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

25 2-(benzyl-sulfonylamino)-4-methyl-pentanoic acid (1-
methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-
3-yl)-amide;

30 2-(beta-styrene-sulfonylamino)-4-methyl-pentanoic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

35 2-((2-methoxycarbonyl)phenyl-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

2-(2-nitro-4-(trifluoromethyl)phenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

5 2-(3-(trifluoromethyl)phenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

10 2-(2,5-dimethoxyphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

15 2-(2-methylphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

20 2-(3,4-dichlorophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

2-(4-(trifluoromethoxy)phenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

25 2-(3,4-dimethoxyphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

30 2-(2-bromophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

35 2-(3,5-bis(trifluoromethyl)phenyl-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

2-(4-ethylphenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

5 2-(4-isopropylphenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

10 2-(2,5-dichlorothiophene-3-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

15 2-(5-chlorothiophene-2-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

20 2-(2-(trifluoromethyl)phenyl-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

25 2-(3-methylphenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

30 2-(2,3-dichlorothiophene-5-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

35 2-(4-bromo-5-chlorothiophene-2-sulfonylamino)-4-
methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-
1H-benzo[e][1,4]diazepin-3-yl)-amide;

35 2-(5-[(benzoylamino)methyl]-thiophene-2-
sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-
phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

2-(4-phenylsulfonylthiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

5 2-(5-(phenylsulfonyl)thiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

10 2-[2-(1-methyl-(5-trifluoromethyl)pyrazole)thiophene-5-sulfonylamino]-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

15 2-(5-(2-pyridyl)thiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

20 2-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-4-methyl-pentanoic acid [5-(4-fluoro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-amide;

N-[1-(3'-Chloro-4'-fluoro-biphenyl-3-ylmethyl)-2-oxo-azepan-3-yl]-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-propionamide;

25 2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-[5-(4-fluoro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-propionamide;

30 2-(2-fluoro-benzenesulfonylamino)-N-[5-(4-fluoro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-propionamide;

35 N-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-2-(2-trifluoromethyl-benzenesulfonylamino)-propionamide;

2-(2-fluoro-benzenesulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

4-methyl-2-(2-trifluoromethyl-benzenesulfonylamino)-
5 pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

4-methyl-2-(propane-1-sulfonylamino)-pentanoic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
10 benzo[e][1,4]diazepin-3-yl)-amide;

4-methyl-2-(propane-1-sulfonylamino)-pentanoic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

15 2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-[1-
methyl-2-oxo-5-(4-trifluoromethyl-phenyl)-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl]-propionamide;

20 2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-4-methyl-
pentanoic acid (7-chloro-1-methyl-2-oxo-5-phenyl-2,3-
dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

4-methyl-2-(propane-1-sulfonylamino)-pentanoic acid
25 [5-(4-chloro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl]-amide;

2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-pyridin-2-yl-2,3-dihydro-
30 1H-benzo[e][1,4]diazepin-3-yl)-amide;

2-[(3,5-dimethyl-isoxazole-4-sulfonyl)-methyl-amino]-
N-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-propionamide;

35 3-dimethylamino-2-(3,5-dimethyl-isoxazole-4-
sulfonylamino)-N-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-propionamide;

N-[1-methyl-2-oxo-5-(4-trifluoromethyl-phenyl)-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-2-(propane-1-sulfonylamino)-propionamide;

5

N-(7-chloro-1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-propionamide;

10

2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-4-methyl-pentanoic acid [5-(4-chloro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-amide; and

15

2-methanesulfonylamino-N-[1-methyl-2-oxo-5-(4-trifluoromethyl-phenyl)-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-propionamide.

In another embodiment, the present invention provides a pharmaceutical composition comprising a compound of Formula (I) and a pharmaceutically acceptable carrier.

In another embodiment, the present invention provides a method for the treatment of neurological disorders comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Formula (I).

In another embodiment, the present invention provides a method for the treatment of neurological disorders associated with A β amyloid production comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Formula (I).

In another embodiment, the present invention provides a method for the treatment of Alzheimer's Disease

comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Formula (I).

5 In another embodiment, the present invention provides a method for the treatment of Alzheimer's Disease associated with A β amyloid production comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Formula
10 (I).

 In another embodiment, the present invention provides a method for for inhibiting γ -secretase activity comprising administering to a host in need of such inhibition a
15 therapeutically effective amount of a compound of Formula (I) that inhibits γ -secretase activity.

 In another embodiment the present invention provides a compound of Formula (I) for use in therapy of Alzheimer's
20 Disease.

 In another embodiment, the present invention provides for the use of a compound of Formula (I) for the manufacture of a medicament for the treatment of
25 Alzheimer's Disease.

DEFINITIONS

 As used herein, the term "A β " denotes the protein
30 designated A β , β -amyloid peptide, and sometimes β /A4, in the art. A β is an approximately 4.2 kilodalton (kD) protein of about 39 to 43 amino acids found in amyloid plaques, the walls of meningeal and parenchymal arterioles, small arteries, capillaries, and sometimes, venules. The
35 isolation and sequence data for the first 28 amino acids

are described in U.S. Pat. No 4,666,829. The 43 amino acid sequence is:

```

1
Asp  Ala  Glu  Phe  Arg  His  Asp  Ser  Gly  Tyr
11
Glu  Val  His  His  Gln  Lys  Leu  Val  Phe  Phe
21
Ala  Glu  Asp  Val  Gly  Ser  Asn  Lys  Gly  Ala
31
Ile  Ile  Gly  Leu  Met  Val  Gly  Gly  Val  Val
41
Ile  Ala  Thr

```

5 The term "APP", as used herein, refers to the protein known in the art as β amyloid precursor protein. This protein is the precursor for A β and through the activity of "secretase" enzymes, as used herein, it is processed into A β . Differing secretase enzymes, known in the art, have
10 been designated γ secretase, generating the N-terminus of A β , γ secretase cleaving around the 16/17 peptide bond in A β , and " γ secretases", as used herein, generating C-terminal A β fragments ending at position 38, 39, 40, 42, and 43 or generating C-terminal extended precursors which
15 are subsequently truncated to the above polypeptides

 The compounds herein described may have asymmetric centers. Compounds of the present invention containing an asymmetrically substituted atom may be isolated in optically active or racemic forms. It is well known in the
20 art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically active starting materials. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable
25 isomers are contemplated in the present invention. Cis and trans geometric isomers of the compounds of the present

invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. All chiral, diastereomeric, racemic forms and all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomeric form is specifically indicated.

The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced.

When any variable (e.g., R^{1a} , R^2 , R^{13} etc.) occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-3 R^{1a} , then said group may optionally be substituted with up to three R^{1a} groups and R^{1a} at each occurrence is selected independently from the definition of R^{1a} . Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

When a bond to a substituent is shown to cross a bond connecting two atoms in a ring, then such substituent may be bonded to any atom on the ring. When a substituent is listed without indicating the atom via which such substituent is bonded to the rest of the compound of a given formula, then such substituent may be bonded via any atom in such substituent. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

As used herein, "alkyl" or "alkylene" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms; for example, "C₁-C₆ alkyl" denotes alkyl having 1 to 6 carbon atoms. Examples of alkyl include, but

are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, sec-butyl, t-butyl, pentyl, and hexyl. Preferred "alkyl" group is "C₁-C₄ alkyl" wherein methyl, ethyl, n-propyl, i-propyl, n-butyl, and i-butyl, are specifically preferred. As used herein, "C₁-C₃ alkyl", whether a terminal substituent or a alkylene group linking two substituents, is understood to specifically include both branched and straight-chain methyl, ethyl, and propyl.

As used herein, "alkenyl" or "alkenylene" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more unsaturated carbon-carbon bonds which may occur in any stable point along the chain. Examples of "C₂-C₆ alkenyl" include, but are not limited to, ethenyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 3-methyl-2-butenyl, 2-pentenyl, 3-pentenyl, hexenyl, and the like.

As used herein, "alkynyl" or "alkynylene" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more carbon-carbon triple bonds which may occur in any stable point along the chain, such as ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, and the like.

"Alkoxy" or "alkyloxy" represents an alkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge. Examples of alkoxy include, but are not limited to, methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, s-butoxy, t-butoxy, n-pentoxy, and s-pentoxy. Preferred alkoxy groups are methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, s-butoxy, t-butoxy. Similarly, "alkylthio" or "thioalkoxy" is represents an alkyl group as defined above with the indicated number of carbon atoms attached through a sulphur bridge.

"Halo" or "halogen" as used herein refers to fluoro, chloro, bromo, and iodo. Unless otherwise specified, preferred halo is fluoro and chloro. "Counterion" is used to represent a small, negatively charged species such as

chloride, bromide, hydroxide, acetate, sulfate, and the like.

"Haloalkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more halogen (for example $-C_vF_w$ where $v = 1$ to 3 and $w = 1$ to $(2v+1)$). Examples of haloalkyl include, but are not limited to, trifluoromethyl, trichloromethyl, pentafluoroethyl, pentachloroethyl, 2,2,2-trifluoroethyl, 2,2-difluoroethyl, heptafluoropropyl, and heptachloropropyl. "Haloalkoxy" is intended to mean a haloalkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge; for example trifluoromethoxy, pentafluoroethoxy, 2,2,2-trifluoroethoxy, and the like. "Halothioalkoxy" is intended to mean a haloalkyl group as defined above with the indicated number of carbon atoms attached through a sulphur bridge.

"Cycloalkyl" is intended to include saturated ring groups, having the specified number of carbon atoms. For example, "C₃-C₆ cycloalkyl" denotes such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

As used herein, "carbocycle" is intended to mean any stable 3- to 7-membered monocyclic or bicyclic or 7- to 13-membered bicyclic or tricyclic, any of which may be saturated, partially unsaturated, or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, or tetrahydronaphthyl (tetralin). Preferred example of "C₃-C₁₀ carbocycle" or "C₃-C₆ carbocycle" is C₃-C₆ cycloalkyl, specifically cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

As used herein, the term "heterocycle" or "heterocyclic ring" is intended to mean a stable 5- to 7-

membered monocyclic or bicyclic or 7- to 14-membered bicyclic heterocyclic ring which is saturated partially unsaturated or unsaturated (aromatic), and which consists of carbon atoms and 1, 2, 3 or 4 heteroatoms, preferably 1, 2, or 3 heteroatoms, independently selected from the group consisting of N, O and S and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may optionally be oxidized. The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. If specifically noted, a nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1.

Examples of heterocycles include, but are not limited to, 1H-indazole, 2-pyrrolidonyl, 2H,6H-1,5,2-dithiazinyl, 2H-pyrrolyl, 3H-indolyl, 4-piperidonyl, 4aH-carbazole, 4H-quinolizinyll, 6H-1,2,5-thiadiazinyl, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazalonyl, carbazolyl, 4aH-carbazolyl, b-carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolinyl, imidazolyl, 1H-indazolyl, indolenyl, indolinyl, indolizinyll, indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl, isoxazolinyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl,


oxazolidinyl, oxazolyl, oxazolidinylperimidinyl,
phenanthridinyl, phenanthrolinyl, phenarsazinyl,
phenazinyl, phenothiazinyl, phenoxathiinyl, phenoxazinyl,
phthalazinyl, piperazinyl, piperidinyl, pteridinyl,
5 piperidonyl, 4-piperidonyl, pteridinyl, purinyl, pyranyl,
pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl,
pyridazinyl, pyridooxazole, pyridoimidazole,
pyridothiazole, pyridinyl, pyridyl, pyrimidinyl,
pyrrolidinyl, pyrrolinyl, pyrrolyl, quinazolinyl,
10 quinolinyl, 4H-quinolizinyll, quinoxalinyl, quinuclidinyl,
carbolinyl, tetrahydrofuranyl, tetrahydroisoquinolinyl,
tetrahydroquinolinyl, 6H-1,2,5-thiadiazinyl, 1,2,3-
thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl,
1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl,
15 thienothiazolyl, thienooxazolyl, thienoimidazolyl,
thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl,
1,2,5-triazolyl, 1,3,4-triazolyl, xanthenyl. Preferred 5
to 10 membered heterocycles include, but are not limited
to, pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl,
20 thiazolyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl,
isoxazolyl, tetrazolyl, benzofuranyl, benzothiofuranyl,
indolyl, benzimidazolyl, 1H-indazolyl, oxazolidinyl,
isoxazolidinyl, benzotriazolyl, benzisoxazolyl, oxindolyl,
benzoxazolinyl, quinolinyl, and isoquinolinyl. Preferred 5
25 to 6 membered heterocycles include, but are not limited to,
pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl,
thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,
imidazolyl, oxazolyl, isoxazolyl, tetrazolyl; more
preferred 5 to 6 membered heterocycles include, but are not
30 limited to, pyridinyl, pyrimidinyl, triazinyl, furanyl,
thienyl, thiazolyl, piperazinyl, piperidinyl, pyrazolyl,
imidazolyl, and tetrazolyl. Also included are fused ring
and spiro compounds containing, for example, the above
heterocycles.

35 As used herein, the term "aryl", "C₆-C₁₀ aryl" or
aromatic residue, is intended to mean an aromatic moiety
containing the specified number of carbon atoms; for
example phenyl, pyridinyl or naphthyl. Unless otherwise

specified, "aryl" may be unsubstituted or substituted with 0 to 3 groups selected from H, OH, OCH₃, Cl, F, Br, I, CN, NO₂, NH₂, N(CH₃)H, N(CH₃)₂, CF₃, OCF₃, C(=O)CH₃, SCH₃, S(=O)CH₃, S(=O)₂CH₃, CH₃, CH₂CH₃, CO₂H, and CO₂CH₃.

5 The term "amino acid" as used herein, refers to natural, modified or unnatural amino acids of either D- or L-configuration and means an organic compound containing both a basic amino group and an acidic carboxyl group. Natural amino acids residues are Ala, Arg, Asn, Asp, Aze,
 10 Cys, Gln, Glu, Gly, His, Hyp, Ile, Irg Leu, Lys, Met, Orn, Phe, Phe(4-fluoro), Pro, Sar, Ser, Thr, Trp, Tyr, and Val. Roberts and Vellaccio, The Peptides, Vol 5; 341-449 (1983), Academic Press, New York, discloses numerous suitable
 15 unnatural amino acids and is incorporated herein by reference for that purpose.


The phrase "additional lactam carbons", as used herein, is intended to denote the number of optional carbon atoms in the lactam ring B of Formula (I). Formula (Ia):




represents the lactam ring B of Formula (I). Additional lactam carbons are carbons in lactam ring B other than the carbons numbered 2 and 3 in the backbone of the formula.

25 The additional lactam carbons may be optionally replaced by a heteroatom selected from oxygen, nitrogen and sulfur. Lactam ring B contains 1, 2, 3, 4, 5, 6 or 7 optional carbons, wherein one optional carbon may optionally be replaced by a heteroatom, such that the total number of
 30 members of lactam ring B, including atoms numbered 1, 2 and 3 in the backbone, does not exceed 10. It is preferred that the total number of atoms of lactam ring B is 6, 7 or 8; it is more preferred that the total number of atoms of lactam ring B is seven. Examples of lactam ring B include:


35




B1




B2




B3




B4




B5




B6




B7




B8




B9






B10



B11



B12




10

B13

B14



B15



B16

- 5 but are not intended to limit the invention. Preferred examples of lactam ring B are B1, B2, B5, B6, B8, B9, B13, and B16; more preferred examples of lactam ring B are B1, B6, B8, B9, and B13; even more preferred examples of lactam ring B are B1 and B6. Preferred examples of substituent
- 10 R^{10} or R^{11} on lactam ring B are methyl, ethyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, cyclopropylethyl, cyclobutylethyl, cyclopentylethyl, cyclohexylethyl,
- 15 4-F-phenyl, 4-Cl-phenyl, 4-CH₃-phenyl, 4-CF₃-phenyl, 4-CH₃O-phenyl, 4-CF₃O-phenyl, 3-F-phenyl, 3-Cl-phenyl, 3-CH₃-phenyl, 3-CF₃-phenyl, 3-CH₃O-phenyl, 3-CF₃O-phenyl, (4-F-phenyl)methyl-, (4-Cl-phenyl)methyl-, (4-CH₃-phenyl)methyl-, (4-CF₃-phenyl)methyl-,
- 20 (4-CH₃O-phenyl)methyl-, (4-CF₃O-phenyl)methyl-, (3-F-phenyl)methyl-, (3-Cl-phenyl)methyl-, (3-CH₃-phenyl)methyl-, (3-CF₃-phenyl)methyl-, (3-CH₃O-phenyl)methyl-, (3-CF₃O-phenyl)methyl-, 2-pyridyl-, 3-pyridyl-, and 4-pyridyl-. More preferred examples of
- 25 substituent R^{10} or R^{11} on lactam ring B are methyl, ethyl, phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-trifluorophenyl, (4-fluorophenyl)methyl, (4-chlorophenyl)methyl, (4-trifluorophenyl)methyl, 2-pyridyl-, 3-pyridyl-, and 4-pyridyl-. The fused rings on lactam ring B may optionally
- 30 be substituted with R^{13} , wherein the preferred examples of substituent R^{13} on fused rings of lactam B are methyl, fluoro, chloro, and methoxy.

The compounds herein described may have asymmetric centers. One enantiomer of a compound of Formula (I) may display superior biological activity over the opposite enantiomer. For example carbon 3 of lactam ring B Formula (I") may exist in either an S or R configuration. Thus, an R or S configuration at carbon 3 in Formula (I") is considered part of the invention. Examples of such configuration include,



but are not intended to be limited to these examples of ring B. When required, separation of the racemic material can be achieved by methods known in the art. Additionally, the carbon atom to which R⁵ is attached may display superior biological activity over the opposite enantiomer. For example, where R⁵ is not H, then the configuration of the carbon may be described as R or S. All configurations are considered part of the invention.

25 The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response,

30

or other problem or complication, commensurate with a reasonable benefit/risk ratio.

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the disclosure of which is hereby incorporated by reference.

"Prodrugs" are intended to include any covalently bonded carriers which release the active parent drug according to Formula (I) *in vivo* when such prodrug is

administered to a mammalian subject. Prodrugs of a compound of Formula (I) are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine
5 manipulation or *in vivo*, to the parent compound. Prodrugs include compounds of Formula (I) wherein a hydroxy, amino, or sulfhydryl group is bonded to any group that, when the prodrug or compound of Formula (I) is administered to a mammalian subject, cleaves to form a free hydroxyl, free
10 amino, or free sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol or acetamide, formamide, benzamide, and N-oxide derivatives of amine functional groups in the compounds of Formula (I), and the
15 like.

"Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic
20 agent.

SYNTHESIS


The compounds of the present invention can be prepared in a number of ways well known to one skilled in the art of organic synthesis. The compounds of the present invention
25 can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or variations thereon as appreciated by those skilled in the art. Preferred methods include, but are not limited to, those described below.
30 All references cited herein are hereby incorporated in their entirety herein by reference.

The novel compounds of this invention may be prepared using the reactions and techniques described in this section. The reactions are performed in solvents
35 appropriate to the reagents and materials employed and are suitable for the transformations being effected. Also, in the description of the synthetic methods described below,

it is to be understood that all proposed reaction conditions, including choice of solvent, reaction atmosphere, reaction temperature, duration of the experiment and workup procedures, are chosen to be the
5 conditions standard for that reaction, which should be readily recognized by one skilled in the art. It is understood by one skilled in the art of organic synthesis that the functionality present on various portions of the molecule must be compatible with the reagents and reactions
10 proposed. Such restrictions to the substituents which are compatible with the reaction conditions will be readily apparent to one skilled in the art and alternate methods must then be used.


15 Compounds of Formula (I) of the present invention can be synthesized by the method of Scheme 1 comprising: step 1, an amino acid coupling; followed by step 2, a deprotection; followed by step 3, a sulfonyl coupling (see Scheme 1). In the method of Scheme 1, a W-X-Y-Z-
20 substituted aminolactam, **X**, is coupled with a protected natural or unnatural amino acid, **XI**, to form a compound, **XII**. The amino protecting group of **XII** is then removed using a standard deprotection procedure to give a compound **XIII**. The compound **XIII** is coupled with an activated
25 sulfonylating agent, **XIV**, preferably a sulfonyl chloride Q-SO₂Cl, **XIVa**, to form a compound of Formula I.

Scheme 1.



P = Amine Protecting group

- 10 In an example of the method of Scheme 1, the coupling of Boc-Alanine to racemic 3-amino-1-methyl-5-phenyl-1,3-dihydro-benzo[e][1,4]diazepin-2-one followed by TFA-mediated removal of the Boc group provides the Alanine-substituted lactam **2b** (Scheme 1a). Sulfonamide formation
- 15 using 3,5-Dimethyl-isoxazole-4-sulfonyl chloride and triethylamine as base provides the compound of Example 2.


Scheme 1a

5

Example 2



Alternatively, compounds of Formula I can be prepared by the method of Scheme 2 which comprises a sequence of sulfonyl coupling, followed by deprotection, followed by amino acid coupling. In the method of Scheme 2, an amino acid derivative which has a protected carboxyl group, **XV**, is first coupled with a sulfonylating agent (Q)SO₂Cl, **XIV**, or an equivalent activated sulfonylating agent, to form a sulfonamide **XVI**. The carboxyl protecting group of **XVI** is removed using a standard deprotection method to give a sulfonamide carboxylic acid **XVII**. The W-X-Y-Z-substituted aminolactam, **X**, is then coupled with the sulfonamide carboxylic acid, **XVII**, to form a compound of Formula I.

Scheme 2.



P* = Carboxylic Acid Protecting Group

5



10 Methods for the synthesis of lactam intermediates as contemplated by the present invention useful in the synthesis of compounds of Formula (I), including amino benzodiazepinones, dibenzo azepinones and other related heterocycles, are known in the art and are disclosed in a
 15 number of references including PCT publication number WO 98/28268, WO 99/66934, WO 00/07995, and WO 00/38618, which are hereby incorporated by reference. Additional references include Bock, et al, J. Org. Chem., 1987, 52, 3232-3239; Sherrill et al, J. Org. Chem., 1995, 60, 730-
 20 734; and Walsh, D. A., Synthesis, September 1980, p.677; and Brown, et al., Tetrahedron Letters, 1971, 8, 667-670.

Synthetic approaches to substituted benzodiazepines are widely described in the literature. The typical methods are illustrated by, but are not limited to, the
 25 following references: M. G. Bock et al J. Org. Chem. 1987, 52, 3232. (b) R. G. Sherrill et al J. Org. Chem. 1995, 60,

734. (c) M. G. Bock et al J. Med. Chem. 1989, 32, 13-16.
(d) J. L. Castro et al J. Med. Chem. 1997, 40, 2491-2501.
(e) M. S. Chambers et al Bioorg. & Med. Chem. Lett. 1993, 3
(10), 1919-1924. (f) J. H. Gogerty et al J. Med. Chem.
5 1977, 20 (7), 952. (g) G. Semple et al Bioorg. & Med. Chem.
Lett. 1996, 6(1), 51-54. (h) G. Semple et al J. Med. Chem.
1997, 40, 331-341. (i) G. Semple et al Bioorg. & Med. Chem.
Lett. 1996, 6 (1), 55-58. (j) G. Semple et al Synth.
Commun. 1996, 26 (4), 721-727. (k) G. A. Showell et al J.
10 Med. Chem. 1994, 37, 719-721. General synthetic
descriptions of 2-aminobenzophenone with various
substitutions used in the preparation of benzodiaepines may
be found in D. A. Walsh Synthesis 1980, 677.


Incorporation of alpha-alkoxy amino acids provides
15 access to additional compounds. alpha-Alkoxy amino acid
derivatives are available from a number of procedures
available in the art. For instance, they may be
synthesized by electrochemical oxidation of malonic acid
half esters in alcoholic solvents as shown in Horikawa,
20 Hiroshi; Iwasaki, Tameo; Matsumoto, Kazuo; Miyoshi, Muneji.
A new synthesis of 2-alkoxy- and 2-acetoxy-2-amino acids by
anodic oxidation. Tetrahedron Lett. 1976, (3), 191-4.
Alternative procedures include alkoxide displacement of
dichloroacetic acid to form an alpha-halo-alpha-
25 alkoxyacetate, followed by aminolysis as reported in Gross,
Hans; Gloede, Joerg; Freiberg, Juergen. alpha-Halo ethers.
XXIX. Glyoxylic acid derivatives from chloromethoxyacetic
acid or dichloroacetic acid. Ann. Chem., Justus Liebig's
(1967), 702 68-74. and similar references.

30 An example of an L- α -amino- β -thio- ϵ -caprolactam, as
shown in Scheme 3, where ring B is the amino lactam of
XVIII and J is a sulfur atom has been reported in the
literature. See S. A. Ahmed et al, FEBS Letters, (1984),
vol. 174, pages 76-9. One skilled in the art can extend
35 this methodology to the synthesis of β -amino and oxygen
containing rings by analogy. The sulfur-containing

molecules can also be oxidized to the sulfoxide and sulfone by methods known to one skilled in the art.

Scheme 3


5



Methods for the alkylation of lactams as contemplated by the present invention in lactam ring B in Formula (I), including amino benzodiazepines and other related heterocycles, are well known in the art. For example, Scheme 4 demonstrates that the lactam nitrogen of compound XIX or other lactam intermediates of the present invention can be alkylated by generating the anion with bases such as LDA, LiHMDS, lithium bis(trimethylsilyl)amide potassium carbonate or sodium hydride in solvents like THF, with or without cosolvents such as DMPU, HMPA or DMF, and reacting this with a variety of groups containing leaving groups (X^{''}) like bromide, iodide, mesylate or tosylate.

Alkylating agents such as α -bromo amides, ketones and acids can be prepared by a number of literature methods including halogenation of amino acids by diazotization or are commercially available. Other suitable alkylating agents such as alkyl, allylic and benzylic halides can be formed form a variety of precursors such as free-radical addition of halides or activation of alcohols, and other chemistries known to those skilled in the art. For discussion of these types of reactions, see Carey, F.A. and Sundberg, R. J., Advanced Organic Chemistry, Part A, New York: Plenum Press, 1990, pages 304-305, 342-347, 695-698.

Scheme 4.



5 A variety of suitably protected natural and unnatural
 amino acid derivatives are commercially available or can be
 prepared by known procedures and can be coupled to the
 aminolactam of Ring B using standard coupling procedures.
 Representative procedures for preparation of amino acids
 10 include the Strecker amino acid synthesis; the Ugi 4-
 component coupling reaction; alkylation of glycine
 benzophenone imine, Seebach's method (reviewed in
 O'Donnell, Martin J.; Fang, Zhiqiang; Seebach's "Self-
 Regeneration of Chirality" and related methods for the
 15 synthesis of α -amino acids, Hecheng Huaxue (1996), 4(4),
 303-316); Schollkopf's bislactam method, and many others.
 For a review, see Duthaler, Rudolf O. Recent developments
 in the stereoselective synthesis of α -amino acids.
 Tetrahedron (1994), 50(6), 1539-650.

20

Protection groups for amine functional group can be
 prepared by methods well known in the literature for amino
 protecting groups as discussed in Theodora W. Greene's book
 "Protective Groups in Organic Synthesis", for example, N-
 25 Boc using di-*t*-butyldicarbonate in an appropriate solvent

like DMSO. Other protecting groups for amino acids include but are not limited to Boc (tert-butyloxycarbonyl), Fmoc (Fluorenylmethyloxycarbonyl), cBz (Benzyloxycarbonyl), and Alloc (Allyloxycarbonyl). Methods for removing these
5 protecting groups are known to those skilled in the art.

Suitable coupling agents include but are not limited to activating agents such as dicyclohexylcarbodiimide (DCC), diisopropylcarbodiimide (DICI), or 1-(3-
10 dimethylaminopropyl)-3-ethylcarbodiimide (EDC) with or without the additive hydroxybenzotriazole (HOBt) or reagents such as HATU (O-(7-azabenzotriazol-1-yl)-1,1,3,3,-tetramethyluronium hexafluorophosphate) or PyBOP (benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium
15 hexafluorophosphate), which are used in a suitable solvent such as dichloromethane or *N,N*-dimethylformamide with a suitable base such as diisopropylethylamine, *N*-methylmorpholine, or triethylamine. Most preferably, the couplings are run using EDC and HOBt as coupling agents
20 with triethylamine as a base in dichloromethane.

Examples

Chemical abbreviations used in the Examples are defined as follows:

- 25 "DMF" for dimethylformamide,
"THF" for tetrahydrofuran,
"DMPU" for 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidone,
"HMPA" for hexamethylphosphoramide,
30 "TBTU" for O-(1H-benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate, and
"BOP" for benzotriazol-1-yloxytris-(dimethylamino)phosphonium hexafluorophosphate.
"LDA" for lithium diisopropyl amide
35 "LiHMDS" for lithium hexamethyl disilazide or lithium bis(trimethylsilyl) amide.
"HPLC" is an abbreviation used herein for high pressure liquid chromatography. Compounds of the present

invention are generally purified by HPLC using conditions known to one skilled in the art. However, unless otherwise indicated, the following conditions are generally applicable. Reverse-phase HPLC can be carried out using a
5 Vydac C-18 column with gradient elution from 10% to 100 % buffer B in buffer A (buffer A: water containing 0.1% trifluoroacetic acid, buffer B: 10% water, 90% acetonitrile containing 0.1% trifluoroacetic acid).
Alternatively, reverse-phase HPLC can be carried out using
10 a Vydac C-18 column with gradient elution from 10% to 90 % acetonitrile in water.

Example 1

2-(3,5-Dimethyl-isoxazole-4-sulfonylamino)-4-methyl-
15 pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide

Step (1a). A solution of 1.2 g (3.46 mmol) of 3-Amino-1-methyl-5-phenyl-1,3-dihydro-benzo[e][1,4]diazepin-2-one and
20 0.818 g (3.53 mmol) of *N*-Boc-(L)-leucine dissolved in 50 mL of dichloromethane at 0°C was treated sequentially with 1.2 mL (8.65 mmol) of triethylamine, 1.06 g (6.92 mmol) of hydroxybenzotriazole, and 0.827 g (4.32 mmol) of EDC. The cooling bath was removed and the solution was stirred for
25 2 h while warming to rt. The reaction solution was then diluted with 20 mL of water and the organic layer was separated and washed with 30 mL of 1 N HCl solution, 30 mL of a saturated NaHCO₃ solution, then dried over Na₂CO₃ and concentrated in vacuo. Chromatography eluting with a
30 gradient of 20 to 50% ethyl acetate in hexanes provided the desired product **1a**. MS (ESI) M+H = 479.4.

Step (1b). The purified material **1a** was dissolved in 50 mL of a 1:1 solution of dichloromethane and trifluoroacetic
35 acid and stirred for 2 h at rt. The solution was directly concentrated and redissolved in 10 mL of toluene. After removal of the solvent 600 mg (35% for 2 steps) of the desired amine was isolated as its TFA salt, MS (ESI) M+H =

379.4. This material was then freebased by dissolving in 20 mL of dichloromethane followed by extracting 3 X with 10 mL of a saturated NaHCO₃ solution, followed by drying the dichloromethane solution with Na₂SO₄ and concentrating to the free amine **1b**.

Step (1c). The free amine **1b** (200 mg, 0.528 mmol) was dissolved in 20 mL of dichloromethane and 147 microliters (1.05 mmol) of triethylamine was added followed by 124 mg (0.63 mmol) of 3,5-dimethyl-isoxazole-4-sulfonyl chloride. After stirring 18 h at rt, 10 mL of water was added and the organic layer was separated. The organic layer was washed with 10 mL of 1 N HCl, 10 mL of a saturated NaHCO₃ solution, and 10 mL of brine. The organic layer was then dried over Na₂SO₄ and concentrated to an oil. Chromatography eluting with a gradient of 20 to 100% ethyl acetate in hexanes allowed the purification of 80 mg (28% yield) of the two diastereomers of the compound of Example 1. MS (ESI) M+H = 538.3).

Example 2

2-(3,5-Dimethyl-isoxazole-4-sulfonylamino)-N-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-propionamide

Step (2a). The compound of Example 2a was synthesized in a manner similar to that of the compound of Example (1a), but using N-Boc-L-Alanine as the amino acid in 50% yield. MS (ESI) M+H = 437.3

Step (2b). The purified material **2a** (1.0 g) was dissolved in 50 mL of a 1:1 solution of dichloromethane and trifluoroacetic acid and stirred for 2 h at rt. The solution was directly concentrated and redissolved in 10 mL of toluene. After removal of the solvent the material was then freebased by dissolving in 20 mL of dichloromethane followed by extracting 3 X with 10 mL of a saturated NaHCO₃

solution. The dichloromethane solution was dried with Na_2SO_4 and concentrated to 0.75 g (96%) of the free amine **2b**. MS (ESI) $M+H = 337.2$).

- 5 **Step (2c).** The free amine **2b** (100 mg, 0.29 mmol) was dissolved in 10 mL of dichloromethane and 83 microliters (0.6 mmol) of triethylamine was added followed by 70 mg (0.36 mmol) of 3,5-dimethyl-isoxazole-4-sulfonyl chloride. After stirring 18 h at rt, 10 mL of water was added and the
10 organic layer was separated. The organic layer was washed with 10 mL of 1 N HCl, 10 mL of a saturated NaHCO_3 solution, and 10 mL of brine. The organic layer was then dried over Na_2SO_4 and concentrated to an oil. Chromatography eluting with a gradient of 20 to 50% ethyl
15 acetate in hexanes allowed the purification of 30 mg (20% yield) of the compound of Example 2. MS (ESI) $M+H = 496.2$.

Example 3

- 20 2-(3-Bromo-5-chloro-thiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide

- The compound of Example 3 was synthesized in a manner similar to that of the compound of Example 1, but using *N*-
25 Boc-L-Alanine as the amino acid in Step (1a) and using 3-bromo-5-chloro-thiophene-2-sulfonyl chloride in Step (1c). MS (ESI) $M+H = 597.0$.

Examples 4-47.

- 30 Additional Examples 4-47, listed below in Table 1, were synthesized by the following procedure: A solution of the free amine **1b** from step (1b) (4.5 mg, 12 micromoles) was dissolved in 0.5 mL of dichloromethane and 3.3 microliters of triethylamine (25 micromoles) was added,
35 followed by 14 micromoles of a corresponding sulfonyl chloride. After stirring for 24 h at room temperature, 100 mg of a mixture of sulfonic acid derived silica gel and

aminopropyl derived silica gel was added with an additional 0.5 mL of dichloromethane. After stirring for 24 h at room temperature, the solution was collected and concentrated. HPLC eluting with a gradient of from 10 to 100% acetonitrile/water provided purified compounds, in a range of from 0.1 to 5 mg each.

Example 4

2-(naphthalene-1-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 569.3.

Example 5

2-(5-dimethylamino naphthalene-1-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 612.4.

Example 6

2-(naphthalene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 569.3.

Example 7

2-(2-acetamido-4-methylthiazole-5-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 597.3.

Example 8

2-(thiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 525.3.

Example 9

2-(quinoline-8-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 570.3.

Example 10

2-(phenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 519.3.

5

Example 11

2-(2,5-dichlorophenyl sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 588.1.

10

Example 12

2-(mesitylene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 561.3.

15

Example 13

2-(3-nitrophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 564.2.

20

Example 14

2-(4-bromophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 598.1.

25

Example 15

2-(4-fluorophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 537.2.

30

Example 16

2-(4-chlorophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 553.7.

35

Example 17

2-(4-acetamidophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 576.3.

Example 18

2-(4-nitrophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 564.2.

Example 19

2-(4-methoxyphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 549.2.

Example 20

2-(4-tert-butylphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 575.3.

Example 21

2-(p-toluene-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 533.2.

Example 22

2-(benzyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 533.2.

Example 23

2-(beta-styrene-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 545.3.

Example 24

2-((2-methoxycarbonyl)phenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 577.3.

Example 25

2-(2-nitro-4-(trifluoromethyl)phenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 633.2.

5

Example 26

2-(3-(trifluoromethyl)phenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 587.2.

10

Example 27

2-(2,5-dimethoxyphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 579.3.

15

Example 28

2-(2-methylphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 533.2.

20

Example 29

2-(3,4-dichlorophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 588.1.

25

Example 30

2-(4-(trifluoromethoxy)phenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 603.2.

30

Example 31

2-(3,4-dimethoxyphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 579.3.

35

Example 32

2-(2-bromophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 598.1.

5

Example 33

2-(3,5-bis(trifluoromethyl)phenyl-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 655.3.

10

Example 34

2-(4-ethylphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 547.3.

15

Example 35

2-(4-isopropylphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 561.3.

20

Example 36

2-(2,5-dichlorothiophene-3-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 594.1.

25

Example 37

2-(5-chlorothiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 559.7.

30

Example 38

2-(2-(trifluoromethyl)phenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 587.2.

35

Example 39

2-(3-methylphenyl-sulfonylamino)-4-methyl-pentanoic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 533.2.

5

Example 40

2-(2,3-dichlorothiophene-5-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 594.1.

10

Example 41

2-(4-bromo-5-chlorothiophene-2-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 639.3.

15

Example 42

2-(5-[(benzoylamino)methyl]-thiophene-2-sulfonylamino)-4-
methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-
1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H =
658.3.

20

Example 43

2-(4-phenylsulfonylthiophene-2-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 665.4.

25

Example 45

2-(5-(phenylsulfonyl)thiophene-2-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 665.4.

30

Example 46

2-[2-(1-methyl-(5-trifluoromethyl)pyrazole)thiophene-5-
sulfonylamino]-4-methyl-pentanoic acid (1-methyl-2-oxo-5-
phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide.


35 MS (ESI) M+H = 673.4.

Example 47

2-(5-(2-pyridyl)thiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 602.4.

- 5 Tables 1 below provides representative Examples of the compounds of Formula (I) of the present invention.

Table 1.




10

Ex. #	R-SO ₂ -	MassSpec (ESI, M+H) Found
3	3-bromo-5-chloro-2-thiophene-SO ₂ -	597.0
4	1-naphthalene-SO ₂ -	569.3
5	5-dimethylamino-naphthalene-1-SO ₂ -	612.4
6	2-naphthalene-SO ₂ -	569.3
7	2-acetamido-4-methyl-5-thiazole-SO ₂ -	597.3
8	2-thiophene-SO ₂ -	525.3
9	8-quinoline-SO ₂ -	570.3
10	phenyl-SO ₂ -	519.2
11	2,5-dichlorophenyl-SO ₂ -	588.1
12	1,3,5-trimethylphenyl-SO ₂ -	561.3
13	3-nitrophenyl-SO ₂ -	564.2
14	4-bromophenyl-SO ₂ -	598.1
15	4-fluorophenyl-SO ₂ -	537.2
16	4-chlorophenyl-SO ₂ -	553.7
17	4-acetamidophenyl-SO ₂ -	576.3
18	4-nitrophenyl-SO ₂ -	564.2
19	4-methoxyphenyl-SO ₂ -	549.2
20	4-tert-butylphenyl-SO ₂ -	575.3
21	4-methylphenyl-SO ₂ -	533.2
22	benzyl-SO ₂ -	533.2
23	beta-styrene-SO ₂ -	545.3
24	2-methoxycarbonylphenyl-SO ₂ -	577.3
25	2-nitro-4-(trifluoromethyl)phenyl-SO ₂ -	632.2
26	3-(trifluoromethyl)phenyl-SO ₂ -	587.2
27	2,5-dimethoxyphenyl-SO ₂ -	579.3
28	2-methylphenyl-SO ₂ -	533.2
29	3,4-dichlorophenyl-SO ₂ -	588.1
30	4-(trifluoromethoxy)phenyl-SO ₂ -	603.2
31	3,4-dimethoxyphenyl-SO ₂ -	579.3
32	2-bromophenyl-SO ₂ -	598.1

33	3,5-bis(trifluoromethyl)phenyl-SO ₂ -	655.3
34	4-ethylphenyl-SO ₂ -	547.3
35	4-isopropylphenyl-SO ₂ -	561.3
36	2,5-dichlorothiophene-3-SO ₂ -	594.1
37	5-chlorothiophene-2-SO ₂ -	559.7
38	2-(trifluoromethyl)phenyl-SO ₂ -	587.2
39	3-methylphenyl-SO ₂ -	533.2
40	2,3-dichlorothiophene-5-SO ₂ -	594.1
41	4-bromo-5-chlorothiophene-2-SO ₂ -	639.3
42	5-[(benzoylamino)methyl]-2-SO ₂ -	658.3
43	4-phenylsulfonylthiophene-2-SO ₂ -	665.4
45	5-(phenylsulfonyl)thiophene-2-SO ₂ -	665.4
46	2(1-methyl-(5-trifluoromethyl)pyrazole)thiophene-5-SO ₂ -	673.4
47	5-(2-pyridyl)thiophene-2-SO ₂ -	602.4

Example 48

- 1-(3,5-Dimethyl-isoxazole-4-sulfonylamino)-
 5 cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide




- 10 **Step (48a)** 1-Aminocyclohexanecarboxylic acid (1.5 grams, 10.5 mmol) was dissolved in 12.5 mL of a 1N solution of sodium hydroxide, 6 mL of water, and 6 mL of tetrahydrofuran. To this solution 3,5-dimethyl-isoxazole-4-sulfonyl chloride (2.45 g, 12.6 mmol) was added and the
 15 solution was stirred at rt for 16 hours. A sufficient portion of 1N NaOH was then added to return the solution to a pH of 9 or higher, and the reaction solution was extracted 2X with a 25 mL portion of dichloromethane and the combined organic layers were discarded. The aqueous
 20 layer was then acidified with 1N HCl until the pH was 3 or lower, and the aqueous layer was extracted 3X with dichloromethane. The organic layers were dried with

magnesium sulfate and concentrated to a white solid **48a** which was used without further purification (234 mg, 7.4%).

Step (48b) The crude acid **48a** (199 mg, 0.66 mmol) was dissolved in 8 mL of CH₂Cl₂ and treated with 0.25 mL (1.5 mmol) of diisopropylethylamine. The reaction solution was cooled to 0 °C and stirred for 5 min. To the reaction mixture was added 1-hydroxybenzotriazole (162 mg, 1.2 mmol) and HATU (285 mg, 0.75 mmol), and this mixture was stirred for 5 min. A 300 mg (0.60 mmol) portion of 3-Amino-1-methyl-5-phenyl-1,3-dihydro-benzo[e][1,4]diazepin-2-one was then added and the reaction solution was allowed to warm to rt and stirred for 16 h. The reaction was quenched with 10 mL of water, and the layers were separated. The organic layer was rinsed with 10% citric acid, 1N Na OH, brine and then concentrated. The resulting oil was dissolved in ethyl acetate, washed with water (6 X 50 mL) and dried over sodium sulfate. After filtration and concentration the compound of Example 48 was obtained as a pale yellow solid **48** (125 mg, 38%), MS (API, MS (M+H)⁺ = 551.

Example 49

N-[1-(3'-Chloro-4'-fluoro-biphenyl-3-ylmethyl)-2-oxo-azepan-3-yl]-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-propionamide



Step (49a) Di-*tert*-butyldicarbonate (10.2 g, 46.7 mmol) was added portion wise to a solution of L-(-)- α -amino- ϵ -caprolactam (5.0 g, 39.0 mmol) in dimethyl sulfoxide (30 mL). After 5 h at rt, the reaction was partitioned between water (100 mL) and ethyl acetate. The combined organic extracts were washed successively with 1 M HCl (50 mL), brine, and dried (MgSO₄) and concentrated *in vacuo*. The residue was recrystallized in 1:1 v/v ether-hexanes, 2

crops yielded the desired product (6.26 g, 70%) as white solid 49a. MS (M+H-BOC)⁺ = 129.

5 **Step (49b)** Boc-caprolactam 49a (5.0 g , 21.9 mmol) was dissolved in 60 mL of THF and chilled to -78 °C. To the chilled solution was added 24 mL of a 1.0 M solution of lithium bis(trimethylsilyl)amide in THF, and the solution was brought to 0 °C and stirred for 15 min. To the anion solution was added 6.5 g (22 mmol) of 3-iodobenzyl bromide
10 (Aldrich) and the the solution was allowed to warm to rt and stirred for 18 h. The reaction solution was diluted with 50 mL of water and extracted 3 x with ethyl acetate. The combined organic layers were dried over MgSO₄ and concentrated in vacuo. The crude product was purified by
15 chromatography eluting with a gradient of 5-20% ethyl acetate/hexanes to afford 7.0 g (72%) of the alkylated Boc-protected caprolactam as a white solid 49b. MS (M+Na)⁺ = 467.

20 **Step (49c)** The caprolactam 49b (1.1 g, 2.47 mmol) was dissolved in 20 mL of a 50% solution of trifluoroacetic acid in dichloromethane. After 2 h at rt, the solution was

Fluoro-3-chlorophenylboronic acid (304 mg, 1.5 mmol) was then added followed by tris(dibenzylideneacetone) dipalladium (180 mg, 0.17 mmol), triphenylphosphine (184 mg, 0.7 mmol) and 1 mL of a 1M solution of potassium carbonate. The reaction solution was heated to reflux for 16 h and allowed to cool. The reaction solution was then partitioned between 25 mL of ethyl acetate and 25 mL of brine and the organic layer was removed. The aqueous layer was extracted with 2 additional 10 mL portions of ethyl acetate and the combined organic layers were dried and concentrated to an oil. Chromatography eluting with a gradient of 1:5 to 2:5 ethyl acetate/hexanes provided the desired biaryl caprolactam (500mg, 83%) as a white solid **49d**. MS (ESI, M+Na⁺) = 540.3.

Step (49e) The biaryl caprolactam **49d** (0.2 g, 0.39 mmol) was dissolved in 20 mL of a 50% solution of trifluoroacetic acid in dichloromethane. After 2 h at rt, the solution was concentrated to an oil. The oil was then dissolved in 10 mL of dichloromethane and 108 µL of triethylamine (.772 mmol) and 91 mg of 3,5-dimethylisoxazole-4-sulfonyl chloride were added. The reaction solution was stirred at rt for 18 h, and then diluted with 10 mL of water and extracted with 2 10 mL portions of dichloromethane. The organic layers were dried and concentrated, and the residual oil was purified by chromatography eluting with 5% methanol in dichloromethane to afford 130 mg (63%) of the compound of Example 49 as a white solid **49**. MS (ESI, M+Na⁺) = 623.4.

30

Example 50

2-(3,5-Dimethyl-isoxazole-4-sulfonylamino)-N-[5-(4-fluorophenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-propionamide

35

Step (50a): {1-[5-(4-Fluoro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-ylcarbamoyl]-ethyl}-carbamic acid tert-butyl ester.

The compound of **50a** was synthesized in a procedure analogous to the synthesis of the compound of **1a** using 0.5 g of 3-Amino-5-(4-fluoro-phenyl)-1-methyl-1,3-dihydro-benzo[e][1,4]diazepin-2-one, and N-Boc-(L)-alanine. In 88% yield.

Step (50b): 2-Amino-N-[5-(4-fluoro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-propionamide.

The compound of **50b** was deprotected using a procedure analogous to the synthesis of the compound of **2b** but using the compound from **50a** and was carried forward without purification.

Step (50c): The compound of Example 50 was synthesized from the compound of **50b** in a procedure analogous to the synthesis of the compound of **2b** using 3,5-dimethyl isoxazole-4-sulfonyl chloride in 31% yield. MS (ESI) M+H = 514.2.

Example 51

2-2-(3,5-Dimethyl-isoxazole-4-sulfonylamino)-4-methyl-pentanoic acid [5-(4-fluoro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-amide

Step (51a): {1-[5-(4-Fluoro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-ylcarbamoyl]-3-methyl-butyl}-carbamic acid tert-butyl ester.

The compound of **51a** was synthesized in a procedure analogous to the synthesis of the compound of **1a** using 0.5 g of 3-Amino-5-(4-fluoro-phenyl)-1-methyl-1,3-dihydro-benzo[e][1,4]diazepin-2-one, and (L)-leucine. In 92% yield.

Step (51b): 2-Amino-4-methyl-pentanoic acid [5-(4-fluoro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-amide.

The compound of **51b** was deprotected using a procedure analogous to the synthesis of the compound of **2b** but using the compound from **51a** and was carried forward without purification.

Step (51c): The compound of Example 51 was synthesized from the compound of **50b** in a procedure analogous to the synthesis of the compound of **2b** using 3,5-dimethyl isoxazole-4-sulfonyl chloride in 36% yield.. MS (ESI) M+H = 556.2.

Example 52

2-(2-Fluoro-benzenesulfonylamino)-N-[5-(4-fluoro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-propionamide

Step (52a): [1-(1-Methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-ylcarbamoyl)-ethyl]-carbamic acid tert-butyl ester.

The compound of **52a** was synthesized in a procedure similar to that of the compound of **1a** but using N-Boc-(L)-alanine as the amino acid in 94% yield.

Step (52b): 2-Amino-N-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-propionamide.

The compound of **52b** was deprotected using a procedure analogous to the synthesis of the compound of **2b** but using the compound from **52a** and was carried forward without purification.

Step (52c): The compound of Example 52 was synthesized from the compound of **52b** in a procedure analogous to the synthesis of the compound of **2b** using 2-fluorobenzene sulfonyl chloride in 68% yield. MS (ESI) M+H = 495.2.

Example 53

N-(1-Methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-2-(2-trifluoromethyl-
5 benzenesulfonylamino)-propionamide

The compound of Example 53 was synthesized in a
procedure analogous to the synthesis of the compound of
Example 52 using 2-trifluoromethyl benzene sulfonyl
chloride in 68% yield. MS (ESI) M+H = 545.2.

10

Example 54

2-(2-Fluoro-benzenesulfonylamino)-4-methyl-pentanoic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide

15 The compound of Example 54 was synthesized in a
procedure analogous to the synthesis of the compound of
Example 1 using the intermediate from **1b** and
2-fluorobenzene sulfonyl chloride in 70% yield. MS (ESI)
M+H = 537.2.

20

Example 55

4-Methyl-2-(2-trifluoromethyl-benzenesulfonylamino)-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide

25 The compound of Example 55 was synthesized in a
procedure analogous to the synthesis of the compound of
Example 1 using the intermediate from **1b** and
2-trifluoromethylbenzene sulfonyl chloride in 65% yield..
MS (ESI) M+H = 587.2.

30

Example 56

4-Methyl-2-(propane-1-sulfonylamino)-pentanoic acid (1-
methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-
3-yl)-amide

35 The compound of Example 56 was synthesized in a
procedure analogous to the synthesis of the compound of
Example 1 using the intermediate from **1b** and

1-propanesulfonyl chloride in 16% yield. MS (ESI) M+H = 485.2.

Example 57

- 5 4-Methyl-2-(propane-1-sulfonylamino)-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide

The compound of Example 57 was synthesized in a procedure analogous to the synthesis of the compound of
10 Example 1 using the intermediate from **1b** and 1-butan sulfonyl chloride. The compound was purified by reverse-phase HLC and isolated in 1% yield.. MS (ESI) M+H = 499.2.

Example 58

- 15 2-(3,5-Dimethyl-isoxazole-4-sulfonylamino)-N-[1-methyl-2-oxo-5-(4-trifluoromethyl-phenyl)-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-propionamide

- 20 **Step (58a):** {1-[1-Methyl-2-oxo-5-(4-trifluoromethyl-phenyl)-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-ylcarbamoyl]-ethyl}-carbamic acid tert-butyl ester.

The compound of **58a** was synthesized in a procedure analogous to the synthesis of the compound of **1a** using 0.5 g
25 of 3-Amino-1-methyl-5-(4-trifluoromethyl-phenyl)-1,3-dihydro-benzo[e][1,4]diazepin-2-one, and N-Boc-(L)-alanine. In 75% yield.

- 30 **Step (58b):** 3-Amino-1-methyl-5-(4-trifluoromethyl-phenyl)-1,3-dihydro-benzo[e][1,4]diazepin-2-one.

The compound of **58b** was deprotected using a procedure analogous to the synthesis of the compound of **2b** but using the compound from **58a** and was carried forward without purification.

35

Step (58c): The compound of Example 58 was synthesized from the compound of **58b** in a procedure analogous to the

synthesis of the compound of **2b** using 3,5-dimethyl isoxazole-4-sulfonyl chloride in 50% yield. MS (ESI) M+H = 564.2.

5

Example 59

2-(3,5-Dimethyl-isoxazole-4-sulfonylamino)-4-methyl-pentanoic acid (7-chloro-1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide

10 **Step (59a):** [1-(7-Chloro-1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-ylcarbamoyl)-3-methyl-butyl]-carbamic acid tert-butyl ester.

The compound of **59a** was synthesized in a procedure analogous to the synthesis of the compound of **1a** using 0.4 g
15 of 3-Amino-7-chloro-1-methyl-5-phenyl-1,3-dihydro-benzo[e][1,4]diazepin-2-one, and N-Boc-(L)-valine. In 93% yield.

Step (59b): 2-Amino-4-methyl-pentanoic acid (7-chloro-1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide.
20

The compound of **59b** was deprotected using a procedure analogous to the synthesis of the compound of **2b** but using the compound from **59a** and was carried forward without
25 purification.

Step (59c): The compound of Example 59 was synthesized from the compound of **59b** in a procedure analogous to the synthesis of the compound of **2b** using 3,5-dimethyl
30 isoxazole-4-sulfonyl chloride in 50% yield. MS (ESI) M+H = 572.2.

Example 60

4-Methyl-2-(propane-1-sulfonylamino)-pentanoic acid [5-(4-chloro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-amide
35

Step (60a): {1-[5-(4-Chloro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-ylcarbamoyl]-3-methyl-butyl}-carbamic acid tert-butyl ester.

The compound of **60a** was synthesized in a procedure analogous to the synthesis of the compound of **1a** using 15.0 g of 3-Amino-5-(4-chloro-phenyl)-1-methyl-1,3-dihydro-benzo[e][1,4]diazepin-2-one and *N*-Boc-(L)-valine. in 53% yield.

Step (60b): 2-Amino-4-methyl-pentanoic acid [5-(4-chloro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-amide.

The compound of **60b** was deprotected using a procedure analogous to the synthesis of the compound of **2b** but using the compound from **60a** and was carried forward without purification.

Step (60c): The compound of Example 60 was synthesized from the compound of **60b** in a procedure analogous to the synthesis of the compound of **2b** using 1-propanesulfonyl chloride in 80% yield. MS (ESI) $M+H = 519.2$.

Example 61

2-(3,5-Dimethyl-isoxazole-4-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-pyridin-2-yl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide

Step (61a): [3-Methyl-1-(1-methyl-2-oxo-5-pyridin-2-yl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-ylcarbamoyl)-butyl]-carbamic acid tert-butyl ester.

The compound of Example **61a** was synthesized in a procedure analogous to the synthesis of the compound of Example **1a** using 0.31 g of 3-Amino-1-methyl-5-pyridin-2-yl-1,3-dihydro-benzo[e][1,4]diazepin-2-one and *N*-Boc-(L)-valine. in 50% yield.

Step (61b): 2-Amino-4-methyl-pentanoic acid (1-methyl-2-oxo-5-pyridin-2-yl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide.

5 The compound of Example **61b** was deprotected using a procedure analogous to the synthesis of the compound of Example **2b** but using the compound from **61a** and was carried forward without purification.

Step (61c): The compound of Example 61 was synthesized from the compound of Example **61b** in a procedure analogous to the synthesis of the compound of Example **2b** using 3,5-dimethyl isoxazole-4-sulfonyl chloride in 39% yield. MS (ESI) M+H = 539.2.

15 **Example 62**

2-[(3,5-Dimethyl-isoxazole-4-sulfonyl)-methyl-amino]-N-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-propionamide

20 The compound from Example 60 (0.02 g) was dissolved in methanol and treated with excess trimethylsilyldiazomethane until a yellow color persisted. The residual reagent was destroyed with acetic acid and the solution was concentrated to provide the compound of Example 63 in 100% yield. MS (ESI) M+H = 510.2.

25

Example 63

3-Dimethylamino-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-propionamide

30

Step (63a): [2-Chloro-1-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-ylcarbamoyl)-ethyl]-carbamic acid tert-butyl ester.

35 The compound of **63a** was synthesized in a procedure analogous to the synthesis of the compound of **1a** using 0.26 g of 3-Amino-1-methyl-5-phenyl-1,3-dihydro-

benzo[e][1,4]diazepin-2-one and *N*-Boc-(*L*)-α-chloroalanine in 11% yield. MS (ESI) *M*+Na = 457.3.

Step (63b): [2-Dimethylamino-1-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-ylcarbamoyl)-ethyl]-carbamic acid tert-butyl ester.

The compound from **63a** (50 mg, 0.1 mmol) was dissolved in 5 mL of acetonitrile and treated with 0.5 mL of 40% dimethylamine in water. The reaction solution was stirred at rt for 2 h and concentrated to provide the compound of **63b** in 100% yield. MS (ESI) *M*+H = 480.3.

Step (63c): 2-Amino-3-dimethylamino-*N*-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-propionamide.

The compound of **63c** was deprotected using a procedure analogous to the synthesis of the compound of **2b** but using the compound from **63b** and was carried forward without purification.

Step (63d): The compound of Example 63 was synthesized from the compound of **61c** in a procedure analogous to the synthesis of the compound of **2b** using 3,5-dimethyl isoxazole-4-sulfonyl chloride in 50% yield. MS (ESI) *M*+H = 539.2.

Example 64

N-[1-Methyl-2-oxo-5-(4-trifluoromethyl-phenyl)-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-2-(propane-1-sulfonylamino)-propionamide

The compound of Example 64 was synthesized from 0.1 g of the compound of **58b** in a procedure analogous to the synthesis of the compound of **2b** using 1-propanesulfonyl chloride in 50% yield. MS (ESI) *M*+H = 511.2.

Example 65

N-(7-Chloro-1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-propionamide

5 The compound of Example 65 was synthesized using a procedure analogous to the synthesis of the compound of Example 59 but using *N*-Boc-(L)-alanine in the second step. MS (ESI) M+H = 530.1.

Example 66

10 2-(3,5-Dimethyl-isoxazole-4-sulfonylamino)-4-methyl-pentanoic acid [5-(4-chloro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-amide

The compound of Example 66 was synthesized from 10 g of the compound of **60b** in a procedure analogous to the
15 synthesis of the compound of **2b** using 3,5-dimethyl isoxazole-4-sulfonyl chloride in 56% yield. MS (ESI) M+H = 572.2.


Example 67

20 2-Methanesulfonylamino-N-[1-methyl-2-oxo-5-(4-trifluoromethyl-phenyl)-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-propionamide

The compound of Example 67 was synthesized from 0.08 g of the compound of **60b** in a procedure analogous to the
25 synthesis of the compound of **58b** using methanesulfonyl chloride in 60% yield. MS (ESI) M+H = 483.1.

Table 2 below provides representative Examples of the compounds of Formula (I) of the present invention.

Table 2.



Ex. #	R-SO ₂ -	R ₅	R ₁₁	R ₁₃
50	3,5-dimethyl-isoxazole-4-SO ₂ -	methyl	4-F-phenyl	H
51	3,5-dimethyl-isoxazole-4-SO ₂ -	i-butyl	4-F-phenyl	H
52	2-fluorophenyl-SO ₂ -	methyl	4-F-phenyl	H
53	2-trifluoromethyl-phenyl-SO ₂ -	methyl	phenyl	H
54	2-fluorophenyl-SO ₂ -	i-butyl	phenyl	H
55	2-trifluoromethyl-phenyl-SO ₂ -	i-butyl	phenyl	H
56	n-propyl-SO ₂ -	i-butyl	phenyl	H
57	n-butyl-SO ₂ -	methyl	4-CF ₃ -phenyl	H
58	3,5-dimethyl-isoxazole-4-SO ₂ -	methyl	4-CF ₃ -phenyl	H
59	3,5-dimethyl-isoxazole-4-SO ₂ -	i-butyl	phenyl	Cl
60	n-propyl-SO ₂ -	i-butyl	4-Cl-phenyl	H
61	3,5-dimethyl-isoxazole-4-SO ₂ -	i-butyl	pyridin-2-yl	H
62	3,5-dimethyl-isoxazole-4-SO ₂ -	methyl	phenyl	H
63	3,5-dimethyl-isoxazole-4-SO ₂ -	(Me) ₂ N-CH ₂ -	phenyl	H
64	n-propyl-SO ₂ -	methyl	4-CF ₃ -phenyl	H
65	3,5-dimethyl-isoxazole-4-SO ₂ -	methyl	phenyl	Cl
66	3,5-dimethyl-isoxazole-4-SO ₂ -	i-butyl	4-Cl-phenyl	H
67	Methyl-SO ₂ -	methyl	4-CF ₃ -phenyl	H

5

Examples 68-99: Examples 68-99 were synthesized in a manner similar to the synthesis of the compounds of Examples 4-47, but using 1-Amino-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide in a yield after RP-HPLC of 0.5 - 5 mg each.

The intermediate, 1-Amino-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide, was synthesized by deprotection of [1-(1-Methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-ylcarbamoyl)-cyclohexyl]-carbamic acid tert-butyl ester in a manner similar to the synthesis of the compound of **1b**.

The intermediate, [1-(1-Methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-ylcarbamoyl)-cyclohexyl]-carbamic acid tert-butyl ester, was synthesized in a manner

similar to the synthesis of the compound of 1a except using 1.5 g (10.5 mmol) of *N*-Boc-1-aminocyclohexane.

Example 68

- 5 1-(naphthalene-1-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 581.7.

Example 69

- 10 1-(naphthalene-2-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 581.7.

Example 70

- 15 1-(thiophene-2-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 537.7.

Example 71

- 20 1-(phenyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 531.6.

Example 72

- 25 1-(2,5-dichlorophenyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 600.5.

Example 73

- 30 1-(2,4,6-trimethylphenyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 573.7.

Example 74

- 35 1-(3-nitrophenyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 576.6.

Example 75

1-(4-bromophenyl-sulfonylamino)-cyclohexanecarboxylic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
5 benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 610.5.

Example 76

1-(4-fluorophenyl-sulfonylamino)-cyclohexanecarboxylic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
10 benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 549.6.

Example 77

1-(4-chlorophenyl-sulfonylamino)-cyclohexanecarboxylic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
15 benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 566.1.

Example 78

1-(beta-styrene-sulfonylamino)-cyclohexanecarboxylic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
20 benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 557.7.

Example 79

1-(4-nitrophenyl-sulfonylamino)-cyclohexanecarboxylic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
25 benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 576.6.

Example 80

1-(4-tert-butylphenyl-sulfonylamino)-cyclohexanecarboxylic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
30 benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 587.7.

Example 81

1-(p-toluene-sulfonylamino)-cyclohexanecarboxylic acid (1-
methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-
35 3-yl)-amide. MS (ESI) M+H = 545.7.

Example 82

1-(benzyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 545.7.

5

Example 83

1-(2-methoxycarbonylphenyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 589.7.

10

Example 84

1-(2-nitro-4-(trifluoromethyl)phenyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 644.6.

15

Example 85

1-(3-(trifluoromethyl)phenyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 599.6.

20

Example 86

1-(2,5-dimethoxyphenyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 591.7.

25

Example 87

1-(2-methoxy-5chloro-1phenyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 596.1.

30

Example 88

1-(3,4-dichlorophenyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 600.5.

35

Example 89

1- (5- ((benzoylamino)methyl)-thiophene-2-sulfonylamino)-
cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-
5 dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H
= 670.8.

Example 90

1- (4- (phenylsulfonyl)thiophene-2-sulfonylamino)-
10 cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-
dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H
= 677.8.

Example 91

15 1- (1-methyl-5- (trifluoromethyl)-imidazole-3-sulfonylamino)-
cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-
dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H
= 685.7.

Example 92

20 1- (4-phenyl-phenyl-sulfonylamino)-cyclohexanecarboxylic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 607.7.

Example 93

25 1- (dibenzofuran-2-sulfonylamino)-cyclohexanecarboxylic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 621.7.

Example 94

30 1- (4-n-butylphenyl-sulfonylamino)-cyclohexanecarboxylic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 587.7.

Example 95

35 1- (2- (2-methylthio- pyrimidin-3-yl)-thiophene-5-
sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-

5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide.
MS (ESI) M+H = 661.8.

Example 96

- 5 1-(4-phenoxy-phenyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 623.7.

Example 97

- 10 1-(3,5-dimethyl-isoxazole-4-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 550.6.

Example 98


- 15 1-(2-(trifluoromethyl)phenyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 599.6.

Example 99

- 20 1-(2-methylphenyl-sulfonylamino)-cyclohexanecarboxylic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide. MS (ESI) M+H = 545.7.
- 25

Table 3 below provides representative Examples of the compounds of Formula (I) of the present invention.

Table 3.



5

Ex. #	R-SO ₂ -	MassSpec (Found)
68	1-naphthalene-SO ₂ -	581.7
69	2-naphthalene-SO ₂ -	581.7
70	2-thiophene-SO ₂ -	537.7
71	phenyl-SO ₂ -	531.6
72	2,5-dichlorophenyl-SO ₂ -	600.5
73	2,4,6-trimethylphenyl-SO ₂ -	573.7
74	3-nitrophenyl-SO ₂ -	576.6
75	4-bromophenyl-SO ₂ -	610.5
76	4-fluorophenyl-SO ₂ -	549.6
77	4-chlorophenyl-SO ₂ -	566.1
78	beta-styrene-SO ₂ -	557.7
79	4-nitrophenyl-SO ₂ -	576.6
80	4-tert-butylphenyl-SO ₂ -	587.7
81	4-methylphenyl-SO ₂ -	545.7
82	benzyl-SO ₂ -	545.7
83	2-methoxycarbonylphenyl-SO ₂ -	589.7
84	2-nitro-4-(trifluoromethyl)phenyl-SO ₂ -	644.6
85	3-(trifluoromethyl)phenyl-SO ₂ -	599.6
86	2,5-dimethoxyphenyl-SO ₂ -	591.7
87	2-methoxy-5-chlorophenyl-SO ₂ -	596.1
88	3,4-dichlorophenyl-SO ₂ -	600.5
89	5-[(benzoylamino)methyl]thiophene-2-SO ₂ -	670.8
90	4-(phenylsulfonyl)thiophene-2-SO ₂ -	677.8
91	1-methyl-5-(trifluoromethyl)imidazole-3-SO ₂ -	685.7
92	4-phenyl-phenyl-SO ₂ -	607.7
93	dibenzofuran-2-SO ₂ -	621.7
94	4-n-butylphenyl-SO ₂ -	587.7
95	2-(2-methylthio-pyrimidin-3-yl)thiophene-5-SO ₂ -	661.8
96	4-phenoxy-phenyl-SO ₂ -	623.7
97	3,5-dimethyl-isoxazole-4-SO ₂ -	550.6
98	2-(trifluoromethyl)phenyl-SO ₂ -	599.6
99	2-methylphenyl-SO ₂ -	545.7

UTILITY

A β production has been implicated in the pathology of Alzheimer's Disease (AD). The compounds of the present invention have utility for the prevention and treatment of AD by inhibiting A β production. Methods of treatment target formation of A β production through the enzymes involved in the proteolytic processing of β amyloid precursor protein. Compounds that inhibit β or γ -secretase activity, either directly or indirectly, control the production of A β . Such inhibition of β or γ -secretases reduces production of A β and is expected to reduce or prevent the neurological disorders associated with A β protein, such as Alzheimer's Disease.

Cellular screening methods for inhibitors of A β production, testing methods for the *in vivo* suppression of A β production, and assays for the detection of secretase activity are known in the art and have been disclosed in numerous publications, including PCT publication number WO 98/22493, EPO publication number 0652009, US patent 5703129 and US patent 5593846; all hereby incorporated by reference.

The compounds of the present invention have utility for the prevention and treatment of disorders involving A β production, such as cerebrovascular disorders.

Compounds of the present invention have been shown to inhibit A β production, as determined by the secretase inhibition assay described below.

Compounds of the present invention have been shown to inhibit A β production, utilizing the C-terminus β amyloid precursor protein accumulation assay described below.

Compounds of Formula (I) are expected to possess γ -secretase inhibitory activity. The γ -secretase inhibitory activity of the compounds of the present invention is

demonstrated using assays for such activity, for example, using the assay described below. Compounds of the present invention have been shown to inhibit the activity of γ -secretase, as determined by the $A\beta$ immunoprecipitation assay.

Compounds provided by this invention should also be useful as standards and reagents in determining the ability of a potential pharmaceutical to inhibit $A\beta$ production. These would be provided in commercial kits comprising a compound of this invention.

As used herein " μ g" denotes microgram, "mg" denotes milligram, "g" denotes gram, " μ L" denotes microliter, "mL" denotes milliliter, "L" denotes liter, "nM" denotes nanomolar, " μ M" denotes micromolar, "mM" denotes millimolar, "M" denotes molar, "nm" denotes nanometer, "SDS" denotes sodium dodecyl sulfate, and "DMSO" denotes dimethyl sulfoxide, and "EDTA" denotes ethylenediaminetetraacetate.

A compound is considered to be active if it has an IC_{50} or K_i value of less than about 100 μ M for the inhibition of $A\beta$ production.

β amyloid precursor protein accumulation assay

A novel assay to evaluate the accumulation of $A\beta$ protein was developed to detect potential inhibitors of secretase. The assay uses the N 9 cell line, characterized for expression of exogenous APP by immunoblotting and immunoprecipitation.

The effect of test compounds on the accumulation of $A\beta$ in the conditioned medium is tested by immunoprecipitation. Briefly, N 9 cells are grown to confluency in 6-well plates and washed twice with 1 x Hank's buffered salt solution. The cells are starved in methionine/cysteine deficient media for 30 min, followed by replacement with fresh deficient media containing 150uCi S^{35} Translabel (Amersham). Test compounds dissolved in DMSO (final concentration 1%) are added together with the addition of

radiolabel. The cells are incubated for 4 h at 37°C in a tissue culture incubator.

At the end of the incubation period, the conditioned medium is harvested and pre-cleared by the addition of 5 μ l
5 normal mouse serum and 50 μ l of protein A Sepharose (Pharmacia), mixed by end-over-end rotation for 30 minutes at 4°C, followed by a brief centrifugation in a microfuge. The supernatant is then harvested and transferred to fresh tubes containing 5 μ g of a monoclonal antibody (clone
10 1101.1; directed against an internal peptide sequence in A β) and 50 μ l protein A Sepharose. After incubation overnight at 4°C, the samples are washed three times with high salt washing buffer (50mM Tris, pH 7.5, 500mM NaCl, 5mM EDTA, 0.5% Nonidet P-40), three times with low salt
15 wash buffer (50mM Tris, pH 7.5, 150mM NaCl, 5mM EDTA, 0.5% Nonidet P-40), and three times with 10mM Tris, pH 7.5. The pellet after the last wash is resuspended in SDS sample buffer (Laemmli, 1970) and boiled for 3 minutes. The supernatant is then fractionated on either 10-20%
20 Tris/Tricine SDS gels or on 16.5% Tris/Tricine SDS gels. The gels are dried and exposed to X-ray film or analyzed by phosphorimaging. The resulting image is analyzed for the presence of A β polypeptides. The steady-state level of A β in the presence of a test compound is compared to wells
25 treated with DMSO (1%) alone. A typical test compound blocks A β accumulation in the conditioned medium, and is therefore considered active, with an IC₅₀ less than 100 μ M.

C-Terminus β Amyloid Precursor Protein Accumulation Assay

30 The effect of test compounds on the accumulation of C-terminal fragments is determined by immunoprecipitation of APP and fragments thereof from cell lysates. N 9 cells are metabolically labeled as above in the presence or absence of test compounds. At the end of the incubation period,
35 the conditioned medium are harvested and cells lysed in RIPA buffer (10 mM Tris, pH 8.0 containing 1% Triton X-100, 1% deoxycholate, 0.1% SDS, 150mM NaCl, 0.125% NaN₃).

Again, lysates are precleared with 5ul normal rabbit serum / 50ul protein A Sepharose, followed by the addition of BC-1 antiserum (15ul;) and 50ul protein A Sepharose for 16 hours at 4°C. The immunoprecipitates are washed as above, bound proteins eluted by boiling in SDS sample buffer and fractionated by Tris/Tricine SDS-PAGE. After exposure to X-ray film or phosphorimager, the resulting images are analyzed for the presence of C-terminal APP fragments. The steady-state level of C-terminal APP fragments is compared to wells treated with DMSO (1%) alone. A typical test compound stimulates C-terminal fragment accumulation in the cell lysates, and is therefore considered active, with an IC₅₀ less than 100 µM.

15 Aβ Immunoprecipitation Assay

This immunoprecipitation assay is specific for γ secretase (i.e., proteolytic activity required to generate the C-terminal end of Aβ either by direct cleavage or generating a C-terminal extended species which is subsequently further proteolyzed). N 9 cells are pulse labeled in the presence of a reported γ secretase inhibitor (MDL 28170) for 1 h, followed by washing to remove radiolabel and MDL 28170. The media is replaced and test compounds are added. The cells are chased for increasing periods of times and Aβ is isolated from the conditioned medium and C-terminal fragments from cell lysates (see above). The test compounds are characterized whether a stabilization of C-terminal fragments is observed and whether Aβ is generated from these accumulated precursor. A typical test compound prevents the generation of Aβ out of accumulated C-terminal fragments and is considered active with an IC₅₀ less than 100 µM.

Dosage and Formulation

35 The compounds of the present invention can be administered orally using any pharmaceutically acceptable dosage form known in the art for such administration. The

active ingredient can be supplied in solid dosage forms such as dry powders, granules, tablets or capsules, or in liquid dosage forms, such as syrups or aqueous suspensions. The active ingredient can be administered alone, but is
5 generally administered with a pharmaceutical carrier. A valuable treatise with respect to pharmaceutical dosage forms is Remington's Pharmaceutical Sciences, Mack Publishing.

The compounds of the present invention can be
10 administered in such oral dosage forms as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. Likewise, they may also be administered in intravenous (bolus or
15 infusion), intraperitoneal, subcutaneous, or intramuscular form, all using dosage forms well known to those of ordinary skill in the pharmaceutical arts. An effective but non-toxic amount of the compound desired can be employed to prevent or treat neurological disorders related
20 to β -amyloid production or accumulation, such as Alzheimer's disease and Down's Syndrome.

The compounds of this invention can be administered by any means that produces contact of the active agent with the agent's site of action in the body of a host, such as a
25 human or a mammal. They can be administered by any conventional means available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or in a combination of therapeutic agents. They can be administered alone, but generally administered with a
30 pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

The dosage regimen for the compounds of the present invention will, of course, vary depending upon known
35 factors, such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration; the species, age, sex, health, medical condition, and weight of the recipient; the nature and extent of the

symptoms; the kind of concurrent treatment; the frequency of treatment; the route of administration, the renal and hepatic function of the patient, and the effect desired. An ordinarily skilled physician or veterinarian can readily
5 determine and prescribe the effective amount of the drug required to prevent, counter, or arrest the progress of the condition.

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily
10 dosage may be administered in divided doses of two, three, or four times daily.

The compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those
15 forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

In the methods of the present invention, the compounds herein described in detail can form the active ingredient, and are typically administered in admixture with suitable pharmaceutical diluents, excipients, or carriers
20 (collectively referred to herein as carrier materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a
30 tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the
35 like; for oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Moreover, when desired or

necessary, suitable binders, lubricants, disintegrating agents, and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or β -lactose, corn
5 sweeteners, natural and synthetic gums such as acacia, tragacanth, or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium
10 acetate, sodium chloride, and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, and the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such
15 as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

Compounds of the present invention may also be coupled
20 with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues.
25 Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone,
30 polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacylates, and crosslinked or amphipathic block copolymers of hydrogels.

Gelatin capsules may contain the active ingredient and powdered carriers, such as lactose, starch, cellulose
35 derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous

release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.


Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance. In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or propyl-paraben, and chlorobutanol.

Suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences, Mack Publishing Company, a standard reference text in this field.

CLAIMS

What is claimed is:

- 5 1. A compound of the Formula (I):



(I)

10 or a pharmaceutically acceptable salt or prodrug thereof,
wherein:

Q is C₁-C₆ alkyl substituted with 0-3 R^{1a};

C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{1b};

15 C₆-C₁₀ aryl substituted with 0-3 R^{1b}; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{1b};

20

R^{1a}, at each occurrence, is independently selected from H,
R^{1b}, Cl, F, Br, I, OR¹⁴, CN, NO₂, =O, NR¹⁹R²⁰, CF₃,
C₁-C₄ alkyl; C₂-C₄ alkenyl; C₂-C₄ alkynyl; C₁-C₄
haloalkyl; C₁-C₄ haloalkoxy;

25

C₃-C₁₀ carbocycle substituted with 0-3 R^{1b};

C₆-C₁₀ aryl substituted with 0-3 R^{1b}; and

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{1b};

30

- R^{1b}, at each occurrence, is independently selected from H, OR¹⁴, Cl, F, Br, I, CN, NO₂, =O, NR¹⁹R²⁰, CF₃, OCF₃, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;
 -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d}, -S(=O)-R^{1d}, -S(=O)₂-R^{1d},
 5 -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d},
 -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d}, -NR^{19b}S(=O)-R^{1d},
 -S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, -OC(=O)-R^{1d};
- C₁-C₆ alkyl substituted with 0-3 R^{1c};
 10 C₂-C₆ alkenyl substituted with 0-2 R^{1c};
 C₂-C₆ alkynyl substituted with 0-2 R^{1c};
 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
 C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};
 C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
 15 5 to 14 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 14 membered
 heterocycle is substituted with 0-3 R^{1f};
- 20 R^{1c}, at each occurrence, is independently selected from H,
 OR¹⁴, Cl, F, Br, I, CN, NO₂, NR¹⁹R²⁰, CF₃,
 C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;
 -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d}, -S(=O)-R^{1d}, -S(=O)₂-R^{1d},
 -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d},
 25 -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d}, -NR^{19b}S(=O)-R^{1d},
 -S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, -OC(=O)-R^{1d};
- C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
 C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};
 30 C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
 5 to 10 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{1f};

R^{1d}, at each occurrence, is independently selected from H,
5 C₁-C₆ alkyl substituted with 0-3 R^{1e};
C₂-C₆ alkenyl substituted with 0-2 R^{1e};
C₂-C₆ alkynyl substituted with 0-2 R^{1e};
C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};
10 C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{1f};

15 R^{1e}, at each occurrence, is independently selected from H,
OR¹⁴, Cl, F, Br, I, CN, NO₂, =O, NR¹⁹R²⁰, CF₃, C₁-C₄
alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkoxy;
20 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};
C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};
C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
25 sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{1f};

R^{1f}, at each occurrence, is independently selected from H,
OR¹⁴, SR¹⁴, Cl, F, Br, I, CN, NO₂, =O, NR¹⁹R²⁰, CF₃,
30 C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkoxy, and C₁-C₄ haloalkyl-S-;

R² is H, C₁-C₆ alkyl, (C₁-C₆ alkoxy) C₁-C₃ alkyl,

C₃-C₆ carbocycle, C₆-C₁₀ aryl, (C₃-C₆ carbocycle)methyl, (C₆-C₁₀ aryl)methyl, (C₆-C₁₀ aryl)ethyl, or 5 to 10 membered heterocycle;

- 5 R⁵ is H, OR¹⁴;
 C₁-C₆ alkyl substituted with 0-3 R^{5b};
 C₁-C₆ alkoxy substituted with 0-3 R^{5b};
 C₂-C₆ alkenyl substituted with 0-3 R^{5b};
 C₂-C₆ alkynyl substituted with 0-3 R^{5b};
 10 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{5c};
 C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
 C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or
 5 to 10 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 15 sulphur, wherein said 5 to 10 membered
 heterocycle is substituted with 0-3 R^{5c};

R^{5a} is H or C₁-C₄ alkyl;

- 20 R^{5b}, at each occurrence, is independently selected from:
 H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
 NR¹⁵R¹⁶; acetyl, -S(C₁-C₄ alkyl), -S(=O)(C₁-C₄
 alkyl), -S(=O)₂(C₁-C₄ alkyl);
 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{5c};
 25 C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
 C₆-C₁₀ aryl substituted with 0-3 R^{5c}; and
 5 to 10 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered
 30 heterocycle is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H,
 OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,

S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁶ is H;

5 C₁-C₆ alkyl substituted with 0-3 R^{6a};
C₃-C₁₀ carbocycle substituted with 0-3 R^{6b}; or
C₆-C₁₀ aryl substituted with 0-3 R^{6b};

R^{6a}, at each occurrence, is independently selected from H,
10 C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
aryl or CF₃;

R^{6b}, at each occurrence; is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, C₁-C₆ alkyl,
15 C₁-C₄ alkoxy, C₁-C₄ haloalkyl, and C₁-C₄ haloalkoxy;

Ring B is a 6, 7, or 8 membered lactam,
wherein the lactam is saturated, partially saturated
or unsaturated;
20 wherein each additional lactam carbon is substituted
with 0-2 R¹¹; and,
optionally, the lactam contains a heteroatom selected
from -N=, -NH-, -N(R¹⁰)-, -O-, -S-, -S(=O)-, and
-S(=O)₂-;

25 additionally, two R¹¹ substituents on adjacent atoms may be
combined to form C₃-C₆ carbocycle fused radical, a
benzo fused radical, or a 5 to 6 membered heteroaryl
fused radical,
30 wherein said 5 to 6 membered heteroaryl fused radical
comprises 1-2 heteroatoms selected from N, O, and S;
wherein said benzo fused radical or 5 to 6 membered
heteroaryl fused radical is substituted with 0-3
R¹³;

35

- R^{10} is H, $C(=O)R^{17}$, $C(=O)OR^{17}$, $C(=O)NR^{18}R^{19}$, $S(=O)_2NR^{18}R^{19}$,
 $S(=O)_2R^{17}$;
C₁-C₆ alkyl substituted with 0-2 R^{10a} ;
C₆-C₁₀ aryl substituted with 0-4 R^{10b} ;
- 5 C₃-C₁₀ carbocycle substituted with 0-3 R^{10b} ; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{10b} ;
- 10 R^{10a} , at each occurrence, is independently selected from H,
C₁-C₆ alkyl, OR^{14} , Cl, F, Br, I, =O, CN, NO₂, $NR^{15}R^{16}$,
CF₃, or aryl substituted with 0-4 R^{10b} ;
- 15 R^{10b} , at each occurrence, is independently selected from H,
OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂,
 $NR^{15}R^{16}$, CF₃, acetyl, SCH₃, $S(=O)CH_3$, $S(=O)_2CH_3$, C₁-C₆
alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄
haloalkoxy, and C₁-C₄ haloalkyl-S-;
- 20 R^{11} , at each occurrence, is independently selected from
H, C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, $NR^{18}R^{19}$,
 $C(=O)R^{17}$, $C(=O)OR^{17}$, $C(=O)NR^{18}R^{19}$, $S(=O)_2NR^{18}R^{19}$,
CF₃;
- 25 C₁-C₆ alkyl substituted with 0-1 R^{11a} ;
C₆-C₁₀ aryl substituted with 0-3 R^{11b} ;
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b} ; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
30 sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{11b} ;

R^{11a}, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
CF₃, or phenyl substituted with 0-3 R^{11b};

5 C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{11b};

10

R^{11b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

15

W is -(CR⁸R^{8a})_p-;

p is 0, 1, 2, 3, or 4;

20 R⁸ and R^{8a}, at each occurrence, are independently selected
from H, F, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl
and C₃-C₈ cycloalkyl;

X is a bond;

25 C₆-C₁₀ aryl substituted with 0-3 R^{Xb};
C₃-C₁₀ cycloalkyl substituted with 0-3 R^{Xb};
C₃-C₁₀ carbocycle substituted with 0-3 R^{Xb}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
30 sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{Xb};

R^{Xb}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,

S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

Y is a bond or -(CR⁹R^{9a})_t-V-(CR⁹R^{9a})_u-;

5

t is 0, 1, 2, or 3;

u is 0, 1, 2, or 3;

10 R⁹ and R^{9a}, at each occurrence, are independently selected from H, F, C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -N(R¹⁹)-, -C(=O)NR^{19b}-, -NR^{19b}C(=O)-, -NR^{19b}S(=O)₂-, -

15 S(=O)₂NR^{19b}-, -NR^{19b}S(=O)-, -S(=O)NR^{19b}-, -C(=O)O-, or -OC(=O)-;

Z is H;

C₁-C₈ alkyl substituted with 0-2 R¹²;

20 C₂-C₆ alkenyl substituted with 0-2 R¹²;

C₂-C₆ alkynyl substituted with 0-2 R¹²;

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or

25 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

R¹² is C₆-C₁₀ aryl substituted with 0-4 R^{12b};

30 C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

R^{12b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
5 haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R¹³, at each occurrence, is independently selected from H,
OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂,
NR¹⁵R¹⁶, or CF₃;

10

R¹⁴, at each occurrence, is independently selected from H,
phenyl, benzyl, C₁-C₆ alkyl, or C₂-C₆ alkoxyalkyl;

15 R¹⁵, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆
alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

R¹⁶, at each occurrence, is independently selected from H,
20 OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-
(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

alternatively, -NR¹⁵R¹⁶ may be a heterocyclic ring selected
from the group piperidinyl, morpholinyl,
25 thiomorpholinyl, pyrrolidinyl, homopiperidinyl,
piperazinyl, and N-methylpiperizinyl;

R¹⁷ is H, aryl, aryl-CH₂-, C₁-C₆ alkyl, or C₂-C₆
alkoxyalkyl;

30

R¹⁸, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆
alkyl) and -S(=O)₂-(C₁-C₆ alkyl);


R¹⁹, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

5 R^{19b}, at each occurrence, is independently selected from H, C₁-C₄ alkyl, phenyl, benzyl, and phenethyl; and

R²⁰ and R²¹, at each occurrence, are independently selected from H, C₁-C₄ alkyl, aryl, and aryl(C₁-C₂ alkyl)-;

10

provided that when Q is phenyl, 3-F-phenyl, benzyl, propyl, or octyl; R² is H; R⁵ is methyl, and -WXYZ is methyl, then Ring B is not



15

; wherein R¹³ is H.

2. A compound of Claim 1, or a pharmaceutically acceptable salt or prodrug thereof, wherein:

20

Q is C₁-C₆ alkyl substituted with 0-3 R^{1a};

C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{1b};

C₆-C₁₀ aryl substituted with 0-3 R^{1b}; or

25

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{1b};

30 R^{1a}, at each occurrence, is independently selected from H, R^{1b}, Cl, F, Br, NR¹⁹R²⁰, CF₃, C₁-C₄ alkyl,

C₁-C₄ haloalkyl;

C₃-C₁₀ carbocycle substituted with 0-3 R^{1b};

C₆-C₁₀ aryl substituted with 0-3 R^{1b}; and

5 to 10 membered heterocycle containing 1 to 4

5 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{1b};

R^{1b}, at each occurrence, is independently selected from

10 H, OR¹⁴, Cl, F, Br, NO₂, NR¹⁹R²⁰, CF₃, OCF₃,

C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;

-C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d}, -S(=O)-R^{1d}, -S(=O)₂-R^{1d},

-N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d},

-NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d}, -NR^{19b}S(=O)-R^{1d},

15 -S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, -OC(=O)-R^{1d};

C₁-C₆ alkyl substituted with 0-3 R^{1c};

C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};

C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and

5 to 14 membered heterocycle containing 1 to 4

20 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 14 membered heterocycle is substituted with 0-3 R^{1f};

R^{1c}, at each occurrence, is independently selected from

25 H, OR¹⁴, Cl, F, Br, NO₂, NR¹⁹R²⁰, CF₃, C₁-C₄ alkoxy,

C₁-C₄ haloalkoxy; -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d},

-S(=O)-R^{1d}, -S(=O)₂-R^{1d}, -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d},

-NR^{19b}C(=O)-R^{1d}, -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d},

-NR^{19b}S(=O)-R^{1d}, -S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d},

30 -OC(=O)-R^{1d};

C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};

C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};
C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
5 sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{1f};

R^{1d}, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, and C₆-C₁₀ aryl substituted with 0-3 R^{1f};

10

R^{1f}, at each occurrence, is independently selected from
H, OR¹⁴, SR¹⁴, Cl, F, Br, I, CN, NO₂, =O, NR¹⁹R²⁰,
CF₃, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

15

R² is H or C₁-C₆ alkyl;

R⁵ is H, OR¹⁴;

C₁-C₆ alkyl substituted with 0-3 R^{5b};
20 C₁-C₆ alkoxy substituted with 0-3 R^{5b};
C₂-C₆ alkenyl substituted with 0-3 R^{5b}; or
C₂-C₆ alkynyl substituted with 0-3 R^{5b};

R^{5a} is H or C₁-C₄ alkyl;

25

R^{5b}, at each occurrence, is independently selected from:
H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
NR¹⁵R¹⁶; acetyl, S(C₁-C₄ alkyl), S(=O)(C₁-C₄
alkyl), S(=O)₂(C₁-C₄ alkyl),

30 C₃-C₁₀ cycloalkyl substituted with 0-3 R^{5c};
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁶ is H, methyl, or ethyl;

Ring B is a seven membered lactam,
wherein the lactam is saturated, partially saturated
or unsaturated;
wherein each additional lactam carbon is substituted
with 0-2 R¹¹; and,
optionally, the lactam contains a heteroatom selected
from -N=, -NH-, -N(R¹⁰)-, -O-, -S-, -S(=O)-, and
-S(=O)₂-;

additionally, two R¹¹ substituents on adjacent atoms may be
combined to form C₃-C₆ carbocycle fused radical, a
benzo fused radical, or a 5 to 6 membered heteroaryl
fused radical,
wherein said 5 to 6 membered heteroaryl fused radical
comprises 1-2 heteroatoms selected from N, O, and S;
wherein said benzo fused radical or 5 to 6 membered
heteroaryl fused radical is substituted with 0-3
R¹³;

R¹⁰ is H, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹,
S(=O)₂R¹⁷;

C₁-C₆ alkyl substituted with 0-2 R^{10a};

C₆-C₁₀ aryl substituted with 0-4 R^{10b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{10b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
5 heterocycle is substituted with 0-3 R^{10b};

R^{10a}, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
CF₃, or aryl substituted with 0-4 R^{10b};

10

R^{10b}, at each occurrence, is independently selected from H,
OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂,
NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆
alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄
15 haloalkoxy, and C₁-C₄ haloalkyl-S-;

R¹¹, at each occurrence, is independently selected from
H, C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹,
C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹,
20 CF₃;

C₁-C₆ alkyl substituted with 0-1 R^{11a};
C₆-C₁₀ aryl substituted with 0-3 R^{11b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
5 to 10 membered heterocycle containing 1 to 4
25 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{11b};

R^{11a}, at each occurrence, is independently selected from H,
30 C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
CF₃, or phenyl substituted with 0-3 R^{11b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{11b};

5

R^{11b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

10

W is -(CR⁸R^{8a})_p-;

p is 0, 1, or 2;

15 R⁸ and R^{8a}, at each occurrence, are independently selected
from H, F, methyl, and ethyl;

X is a bond;

phenyl substituted with 0-3 R^{Xb};

20

C₃-C₆ cycloalkyl substituted with 0-3 R^{Xb}; or

5 to 6 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 6 membered heterocycle
is substituted with 0-2 R^{Xb};

25

R^{Xb}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

30

Y is a bond or -(CR⁹R^{9a})_t-V-(CR⁹R^{9a})_u-;

t is 0, 1, or 2;

35 u is 0, 1, or 2;

R⁹ and R^{9a}, at each occurrence, are independently selected from H, F, methyl, and ethyl;

5 V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-,
-N(R¹⁹)-, -NHC(=O)-, or -C(=O)NH-;

Z is H;

C₁-C₆ alkyl substituted with 0-2 R¹²;

10 C₂-C₆ alkenyl substituted with 0-2 R¹²;

C₂-C₆ alkynyl substituted with 0-2 R¹²;

phenyl substituted with 0-4 R^{12b};

C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4
15 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b};

R¹² is phenyl substituted with 0-4 R^{12b};

20 C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b};

25

R^{12b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

30

R¹³, at each occurrence, is independently selected from
H, OH, C₁-C₄ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴, at each occurrence, is independently selected from H, phenyl, benzyl, C₁-C₆ alkyl, and C₂-C₆ alkoxyalkyl;

5 R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

10 R¹⁶, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

15 alternatively, -NR¹⁵R¹⁶ may be a heterocyclic ring selected from the group piperidinyl, morpholinyl, thiomorpholinyl, pyrrolidinyl, homopiperidinyl, piperazinyl, and N-methylpiperiziny;


R¹⁷ is H, aryl, aryl-CH₂-, C₁-C₆ alkyl, or C₂-C₆ alkoxyalkyl;

20 R¹⁸, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

25 R¹⁹, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl); and




R²⁰ is H or C₁-C₄;

30 provided that when Q is phenyl, 3-F-phenyl, benzyl, propyl, or octyl; R² is H; R⁵ is methyl, and -WXYZ is methyl, then Ring B is not




3. A compound of Claim 2 wherein Ring B is selected from:

5




10



wherein each benzo fused radical is substituted with 0-3 R^{13} .

4. A compound of Claim 3 of Formula (I):



5

(I)

or a pharmaceutically acceptable salt form or prodrug thereof, wherein:

Q is C₁-C₆ alkyl substituted with 0-3 R^{1a} ;

10

C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1b} ;

C₆-C₁₀ aryl substituted with 0-3 R^{1b} ; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered

15

heterocycle is substituted with 0-3 R^{1b} ;

R^{1a} , at each occurrence, is independently selected from H, Cl, F, Br, $NR^{19}R^{20}$, CF_3 , C₁-C₄ alkyl; C₁-C₄ haloalkyl;

20

C₆-C₁₀ aryl substituted with 0-3 R^{1b} ; and

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{1b} ;

25

R^{1b} , at each occurrence, is independently selected from H, OR^{14} , Cl, F, Br, NO_2 , $NR^{19}R^{20}$, CF_3 , OCF_3 , C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;

30

-C(=O)- R^{1d} , -O- R^{1d} , -S- R^{1d} , -S(=O)- R^{1d} , -S(=O)₂- R^{1d} ,
-N(R^{19})- R^{1d} , -C(=O)N R^{19b} - R^{1d} , -N R^{19b} C(=O)- R^{1d} ,
-N R^{19b} S(=O)₂- R^{1d} , -S(=O)₂N R^{19b} - R^{1d} , -N R^{19b} S(=O)- R^{1d} ,

-S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, -OC(=O)-R^{1d};

C₁-C₆ alkyl substituted with 0-3 R^{1c};

C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};

5 C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and

5 to 14 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 14 membered
heterocycle is substituted with 0-3 R^{1f};

10

R^{1c}, at each occurrence, is independently selected from H,
OR¹⁴, Cl, F, Br, NO₂, NR¹⁹R²⁰, CF₃, C₁-C₄ alkoxy,
C₁-C₄ haloalkoxy; -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d},
-S(=O)-R^{1d}, -S(=O)₂-R^{1d}, -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d},
15 -NR^{19b}C(=O)-R^{1d}, -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d},
-NR^{19b}S(=O)-R^{1d}, -S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d},
-OC(=O)-R^{1d};

C₃-C₁₀ cycloalkyl substituted with 0-3 R^{1f};

20 C₃-C₁₀ carbocycle substituted with 0-3 R^{1f};

C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered

25 heterocycle is substituted with 0-3 R^{1f};

R^{1d}, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, and

C₆-C₁₀ aryl substituted with 0-3 R^{1f};

30

R^{1f}, at each occurrence, is independently selected from
H, Cl, F, Br, NO₂, CF₃, C₁-C₄ alkyl, C₁-C₄ haloalkyl,
C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, and

C₁-C₄ haloalkyl-S-;

R² is H, methyl, or ethyl;

5 R⁵ is OR¹⁴;

C₁-C₆ alkyl substituted with 0-3 R^{5b};

C₁-C₆ alkoxy substituted with 0-3 R^{5b};

C₂-C₆ alkenyl substituted with 0-3 R^{5b}; or

C₂-C₆ alkynyl substituted with 0-3 R^{5b};

10

R^{5a} is H or C₁-C₄ alkyl;

R^{5b}, at each occurrence, is independently selected from:

H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,

15

NR¹⁵R¹⁶, acetyl, -S(C₁-C₄ alkyl), S(=O)(C₁-C₄ alkyl), S(=O)₂(C₁-C₄ alkyl);

C₃-C₁₀ cycloalkyl substituted with 0-3 R^{5c};

C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or

5 to 10 membered heterocycle containing 1 to 4

20

heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H,


25

OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁶ is H;

30

Ring B is selected from:



R¹¹, at each occurrence, is independently selected from
 H, C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹,
 5 C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹,
 CF₃;

C₁-C₆ alkyl substituted with 0-1 R^{11a};

C₆-C₁₀ aryl substituted with 0-3 R^{11b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or

10 5 to 10 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered
 heterocycle is substituted with 0-3 R^{11b};

15 R^{11a}, at each occurrence, is independently selected from H,
 C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
 CF₃, phenyl substituted with 0-3 R^{11b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or

20 5 to 10 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered heterocycle
 is substituted with 0-3 R^{11b};

R^{11b}, at each occurrence, is independently selected from H,
 25 OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
 S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
 haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

W is -(CR⁸R^{8a})_p-;

p is 0, 1, or 2;

R⁸ and R^{8a}, at each occurrence, are independently selected from H, F, methyl, and ethyl;

5

X is a bond;

phenyl substituted with 0-3 R^{Xb};

C₃-C₆ cycloalkyl substituted with 0-3 R^{Xb}; or

5 to 6 membered heterocycle containing 1 to 4

10 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-2 R^{Xb};

R^{Xb}, at each occurrence, is independently selected from H,
15 OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

Y is a bond or -(CR⁹R^{9a})_t-V-(CR⁹R^{9a})_u-;

20

t is 0, 1, or 2;

u is 0, 1, or 2;

25 R⁹ and R^{9a}, at each occurrence, are independently selected from H, F, methyl, and ethyl;

V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-,
-N(R¹⁹)-, -C(=O)NH-, or -NHC(=O)-;

30

Z is H;

C₁-C₆ alkyl substituted with 0-2 R¹²;

C₂-C₆ alkenyl substituted with 0-2 R¹²;

C₂-C₆ alkynyl substituted with 0-2 R¹²;

35 phenyl substituted with 0-4 R^{12b};

C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
5 heterocycle is substituted with 0-3 R^{12b};

R¹² is phenyl substituted with 0-4 R^{12b};
C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4
10 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b};

R^{12b}, at each occurrence, is independently selected from H,
15 OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R¹³, at each occurrence, is independently selected from
20 H, OH, C₁-C₄ alkyl, C₁-C₄ alkoxy, Cl, F, Br, CN, NO₂,
and CF₃;

R¹⁴, at each occurrence, is independently selected from H,
phenyl, benzyl, C₁-C₆ alkyl, or C₂-C₆ alkoxyalkyl;
25

R¹⁵, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆
alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

30 R¹⁶, at each occurrence, is independently selected from H,
OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-
(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

alternatively, $-NR^{15}R^{16}$ may be a heterocyclic ring selected from the group piperidinyl, morpholinyl, thiomorpholinyl, pyrrolidinyl, homopiperidinyl, piperazinyl, and N-methylpiperizinyl;

5

R^{17} is H, aryl, aryl- CH_2 -, C_1 - C_6 alkyl, or C_2 - C_6 alkoxyalkyl;

R^{18} , at each occurrence, is independently selected from H, C_1 - C_6 alkyl, phenyl, benzyl, phenethyl, $-C(=O)-(C_1-C_6$ alkyl) and $-S(=O)_2-(C_1-C_6$ alkyl);


R^{19} , at each occurrence, is independently selected from H, OH, C_1 - C_6 alkyl, phenyl, benzyl, phenethyl, $-C(=O)-(C_1-C_6$ alkyl) and $-S(=O)_2-(C_1-C_6$ alkyl); and

15

R^{20} is H or C_1 - C_4 ;


provided that when Q is phenyl, 3-F-phenyl, benzyl, propyl, or octyl; R^2 is H; R^5 is methyl, and -WXYZ is methyl, then Ring B is not

20



25

5. A compound of Claim 4 of Formula (Ia):




(Ia)

or a pharmaceutically acceptable salt form or prodrug thereof, wherein:

30

Ring B is



- 5 Q is C₁-C₆ alkyl substituted with 0-2 R^{1a};
 C₆-C₁₀ aryl substituted with 0-3 R^{1b}; or
 5 to 10 membered heterocycle containing 1 to 3
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered
 10 heterocycle is substituted with 0-3 R^{1b};
- R^{1a}, at each occurrence, is independently selected from
 H, Cl, F, Br, NR¹⁹R²⁰, CF₃, C₁-C₄ alkyl, C₁-C₄
 haloalkyl, and phenyl substituted with 0-3 R^{1b};
- 15 R^{1b}, at each occurrence, is independently selected from H,
 OR¹⁴, Cl, F, Br, NO₂, NR¹⁹R²⁰, CF₃, OCF₃, C₁-C₄
 alkoxy, C₁-C₄ haloalkoxy; -C(=O)-R^{1d}, -O-R^{1d}, -S(=O)₂-
 R^{1d}, -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d},
 20 -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d}, -C(=O)O-R^{1d},
 -OC(=O)-R^{1d};
 C₁-C₆ alkyl substituted with 0-3 R^{1c};
 C₆-C₁₀ aryl substituted with 0-3 R^{1f}; and
 5 to 14 membered heterocycle containing 1 to 4
 25 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 14 membered
 heterocycle is substituted with 0-3 R^{1f};
- R^{1c}, at each occurrence, is independently selected from
 30 H, -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d}, -S(=O)-R^{1d},

-S(=O)₂-R^{1d}, -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d},
 -NR^{19b}C(=O)-R^{1d}, -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d},
 -NR^{19b}S(=O)-R^{1d}, -S(=O)NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, and
 -OC(=O)-R^{1d};

5

R^{1d}, at each occurrence, is independently selected from
 H, C₁-C₆ alkyl, and

C₆-C₁₀ aryl substituted with 0-3 R^{1f};

10 R^{1f}, at each occurrence, is independently selected from
 H, Cl, F, Br, NO₂, CF₃, C₁-C₄ alkyl, C₁-C₄ haloalkyl,
 C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, and
 C₁-C₄ haloalkyl-S-;

15 R⁵ is OR¹⁴;

C₁-C₆ alkyl substituted with 0-1 R^{5b};

C₂-C₆ alkenyl substituted with 0-1 R^{5b}; or

C₂-C₆ alkynyl substituted with 0-1 R^{5b};

20 R^{5a} is H or C₁-C₄ alkyl;

R^{5b}, at each occurrence, is independently selected from:

H, methyl, ethyl, propyl, butyl, CF₃, OR¹⁴, Cl, F, Br,

I, =O, CN, NO₂, NR¹⁵R¹⁶; acetyl, -S(C₁-C₄ alkyl),

25 -S(=O)(C₁-C₄ alkyl), -S(=O)₂(C₁-C₄ alkyl),

C₃-C₇ cycloalkyl substituted with 0-3 R^{5c};

phenyl substituted with 0-3 R^{5c}; and

5 to 7 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

30 sulphur, wherein said 5 to 7 membered heterocycle
 is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from

H, OH, Cl, F, -NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)₂CH₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

- 5 R¹¹, at each occurrence, is independently selected from
H, -NR¹⁸R¹⁹, -C(=O)R¹⁷, -C(=O)OR¹⁷, -C(=O)NR¹⁸R¹⁹,
-S(=O)₂NR¹⁸R¹⁹, -CF₃;
C₁-C₆ alkyl substituted with 0-1 R^{11a};
phenyl substituted with 0-3 R^{11b};
10 C₃-C₆ carbocycle substituted with 0-3 R^{11b}; or
5 to 7 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle
is substituted with 0-3 R^{11b};
15 R^{11a}, at each occurrence, is independently selected from H,
C₁-C₄ alkyl, OR¹⁴, Cl, F, Br, CN, NO₂, NR¹⁵R¹⁶, CF₃,
phenyl substituted with 0-3 R^{11b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
20 5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{11b};
25 R^{11b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

30 W is -(CR⁸R^{8a})_p-;

p is 0, 1, or 2;

R^8 and R^{8a} , at each occurrence, are independently selected from H, F, methyl, and ethyl;

X is a bond;

- 5 phenyl substituted with 0-3 R^{Xb} ;
 C_3 - C_6 cycloalkyl substituted with 0-3 R^{Xb} ; or
 5 to 6 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 6 membered heterocycle
10 is substituted with 0-2 R^{Xb} ;

- R^{Xb} , at each occurrence, is independently selected from H,
 OH, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 ,
 $S(=O)CH_3$, $S(=O)_2CH_3$, C_1 - C_6 alkyl, C_1 - C_4 alkoxy, C_1 - C_4
15 haloalkyl, C_1 - C_4 haloalkoxy, and C_1 - C_4 haloalkyl-S-;

Y is a bond or $-(CR^9R^{9a})_t-V-(CR^9R^{9a})_u-$;

t is 0, 1, or 2;

20

u is 0, 1, or 2;

R^9 and R^{9a} , at each occurrence, are independently selected from H, F, methyl, and ethyl;

25

V is a bond, $-C(=O)-$, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$,
 $-N(R^{19})-$, $-C(=O)NH-$, or $-NHC(=O)-$;

Z is H;

- 30 C_1 - C_6 alkyl substituted with 0-2 R^{12} ;
 C_2 - C_6 alkenyl substituted with 0-2 R^{12} ;
 C_2 - C_6 alkynyl substituted with 0-2 R^{12} ;
 phenyl substituted with 0-4 R^{12b} ;
 C_3 - C_6 carbocycle substituted with 0-4 R^{12b} ; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b};

5

R¹² is phenyl substituted with 0-4 R^{12b};

C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b};

10

R^{12b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

15

R¹³, at each occurrence, is independently selected from
H, OH, C₁-C₃ alkyl, C₁-C₃ alkoxy, Cl, F, Br, CN, NO₂,
and CF₃;

20

R¹⁴, at each occurrence, is independently selected from H,
phenyl, benzyl, C₁-C₆ alkyl, and C₂-C₆ alkoxyalkyl;

25 R¹⁵, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆
alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

R¹⁶, at each occurrence, is independently selected from H,
OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-
(C₁-C₆ alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

30

alternatively, -NR¹⁵R¹⁶ may be a heterocyclic ring selected
from the group piperidinyl, morpholinyl,

thiomorpholinyl, pyrrolidinyl, homopiperidinyl,
piperazinyl, and N-methylpiperizinyl;

5 R^{17} is H, phenyl, benzyl, C₁-C₄ alkyl, or C₂-C₄
alkoxyalkyl;


10 R^{18} , at each occurrence, is independently selected from H,
C₁-C₆ alkyl, phenyl, benzyl, phenethyl, -C(=O)-(C₁-C₆
alkyl) and -S(=O)₂-(C₁-C₆ alkyl);

15 R^{19} , at each occurrence, is independently selected from H,
OH, methyl, ethyl, propyl, butyl, phenyl, benzyl, and
phenethyl; and

20 R^{20} is H, methyl, ethyl, propyl, or butyl.

6. A compound of Claim 5 of Formula (Ia) wherein:

20 Ring B is



Q is C₁-C₆ alkyl substituted with 0-1 R^{1a} ;
phenyl substituted with 0-3 R^{1b} ;
25 naphthyl substituted with 0-3 R^{1b} ; or
5 to 10 membered heterocycle containing 1 to 2
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{1b} ;

30

R^{1a} , at each occurrence, is independently selected from

H, Cl, F, Br, -NR¹⁹R²⁰, -CF₃; and
phenyl substituted with 0-3 R^{1b};

R^{1b}, at each occurrence, is independently selected from
5 H, methyl, ethyl, OR¹⁴, Cl, F, Br, NO₂, NR¹⁹R²⁰, CF₃,
OCF₃, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy;
-C(=O)-R^{1d}, -O-R^{1d}, -S(=O)₂-R^{1d}, -N(R¹⁹)-R^{1d},
-C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d}, -NR^{19b}S(=O)₂-R^{1d},
-S(=O)₂NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, -OC(=O)-R^{1d};
10 C₁-C₆ alkyl substituted with 0-3 R^{1c};
phenyl substituted with 0-3 R^{1f}; and
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
15 heterocycle is substituted with 0-3 R^{1f};

R^{1c}, at each occurrence, is independently selected from
H, -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d}, -S(=O)-R^{1d},
-S(=O)₂-R^{1d}, -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d},
20 -NR^{19b}C(=O)-R^{1d}, -NR^{19b}S(=O)₂-R^{1d}, and
-S(=O)₂NR^{19b}-R^{1d};

R^{1d}, at each occurrence, is independently selected from
H, C₁-C₆ alkyl; and phenyl substituted with 0-3 R^{1f};

25 R^{1f}, at each occurrence, is independently selected from
H, Cl, F, Br, NO₂, CF₃, methyl, ethyl, propyl,
methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and
C₁-C₂ haloalkoxy;

30 R⁵ is OR¹⁴ or C₁-C₄ alkyl substituted with 0-1 R^{5b};

R^{5a} is H;

- R^{5b}, at each occurrence, is independently selected from:
H, methyl, ethyl, propyl, butyl, -NR¹⁵R¹⁶;
C₃-C₇ cycloalkyl substituted with 0-3 R^{5c};
5 phenyl substituted with 0-3 R^{5c}; or
5 to 7 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle
is substituted with 0-3 R^{5c}; wherein said 5 to 7
10 membered heterocycle is selected from pyridinyl,
pyrimidinyl, triazinyl, furanyl, thienyl,
thiazolyl, pyrrolyl, piperazinyl, piperidinyl,
homopiperidinyl, pyrazolyl, imidazolyl, oxazolyl,
isoxazolyl, and tetrazolyl;
15
- R^{5c}, at each occurrence, is independently selected from H,
OH, Cl, F, -NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)₂CH₃,
methyl, and methoxy;
- 20 R¹¹, at each occurrence, is independently selected from
H, NR¹⁸R¹⁹, CF₃;
C₁-C₆ alkyl substituted with 0-1 R^{11a};
phenyl substituted with 0-3 R^{11b};
C₃-C₆ carbocycle substituted with 0-3 R^{11b}; or
25 5 to 7 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle
is substituted with 0-3 R^{11b}; wherein said 5 to 7
membered heterocycle is selected from pyridinyl,
30 pyrimidinyl, triazinyl, furanyl, thienyl,
thiazolyl, pyrrolyl, piperazinyl, piperidinyl,
homopiperidinyl, pyrazolyl, imidazolyl, oxazolyl,
isoxazolyl, and tetrazolyl;

R^{11a}, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, Cl, F, Br, CN, NO₂, NR¹⁵R¹⁶, CF₃, phenyl substituted with 0-3 R^{11b};

5 C₃-C₆ carbocycle substituted with 0-3 R^{11b}; or
5 to 7 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 7 membered heterocycle is substituted with 0-3 R^{11b}; wherein said 5 to 7
10 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, homopiperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

15 R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂
20 haloalkoxy;

W is -(CHR⁸)_p-;

p is 0 or 1;

25 R⁸ is H, methyl, or ethyl;

X is a bond;
phenyl substituted with 0-2 R^{Xb};

30 C₃-C₆ cycloalkyl substituted with 0-3 R^{Xb}; or
5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-1 R^{Xb};

35

R^{Xb}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

5

Y is a bond, -V-, -CH₂-V-, -V-CH₂-, or -CH₂-V-CH₂-;

V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-,
-N(R¹⁹)-, -C(=O)NH-, or -NHC(=O)-;

10

Z is H;

C₁-C₆ alkyl substituted with 0-2 R¹²;

C₂-C₆ alkenyl substituted with 0-2 R¹²;

C₂-C₆ alkynyl substituted with 0-2 R¹²;

15

phenyl substituted with 0-4 R^{12b};

C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered

20

heterocycle is substituted with 0-3 R^{12b};

R¹² is phenyl substituted with 0-4 R^{12b};

C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b};

25

R^{12b}, at each occurrence, is independently selected from H,
30 OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R¹³, at each occurrence, is independently selected from

H, methyl, ethyl, methoxy, ethoxy, Cl, F, Br, NO₂,
or CF₃;

R¹⁴, at each occurrence, is independently selected from H,
5 phenyl, benzyl, C₁-C₄ alkyl, or C₂-C₄ alkoxyalkyl;

R¹⁵, at each occurrence, is independently selected from H,
methyl, ethyl, propyl, butyl, phenyl, benzyl, and
phenethyl;

10

R¹⁶, at each occurrence, is independently selected from H,
OH, methyl, ethyl, propyl, butyl, phenyl, benzyl,
phenethyl, CH₃CH₂C(=O)-, CH₃C(=O)-, CH₃CH₂OC(=O)-,
CH₃OC(=O)-, CH₃CH₂S(=O)₂- and CH₃S(=O)₂-;

15


R¹⁸, at each occurrence, is independently selected from H,
methyl, ethyl, propyl, butyl, phenyl, benzyl, and
phenethyl;

20 R¹⁹, at each occurrence, is independently selected from H,
OH, methyl, ethyl, propyl, and butyl; and

R²⁰ is H, methyl, ethyl, propyl, or butyl.

25 7. A compound of Claim 6 of Formula (Ia) wherein:

Ring B is



30 Q is C₁-C₄ alkyl substituted with 0-1 R^{1a};
phenyl substituted with 0-3 R^{1b};
naphthyl substituted with 0-3 R^{1b}; or

- 5 to 10 membered heterocycle containing 1 to 2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{1b}; wherein said 5 to 10 membered heterocycle is selected from pyridinyl, quinolinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, imidazolyl, oxazolyl, and isoxazolyl;
- 10 R^{1a}, at each occurrence, is independently selected from H, Cl, F, Br, CF₃, and phenyl substituted with 0-3 R^{1b};
- R^{1b}, at each occurrence, is independently selected from H, methyl, ethyl, OR¹⁴, Cl, F, Br, NO₂, NR¹⁹R²⁰, CF₃, OCF₃, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy; -C(=O)-R^{1d}, -O-R^{1d}, -S(=O)₂-R^{1d}, -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d}, -NR^{19b}S(=O)₂-R^{1d}, -S(=O)₂NR^{19b}-R^{1d}, -C(=O)O-R^{1d}, -OC(=O)-R^{1d};
- 20 C₁-C₆ alkyl substituted with 0-3 R^{1c}; phenyl substituted with 0-3 R^{1f}; and 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{1f};
- 25 R^{1c}, at each occurrence, is independently selected from H, -C(=O)-R^{1d}, -O-R^{1d}, -S-R^{1d}, -S(=O)-R^{1d}, -S(=O)₂-R^{1d}, -N(R¹⁹)-R^{1d}, -C(=O)NR^{19b}R^{1d}, -NR^{19b}C(=O)-R^{1d}, -NR^{19b}S(=O)₂-R^{1d}, and -S(=O)₂NR^{19b}-R^{1d};
- 30 R^{1d}, at each occurrence, is independently selected from H, C₁-C₆ alkyl; and phenyl substituted with 0-3 R^{1f};

R^{1f}, at each occurrence, is independently selected from
H, Cl, F, Br, NO₂, CF₃, -OCF₃, methyl, ethyl, methoxy,
and ethoxy;

5

R⁵ is methyl substituted with 0-1 R^{5b};
ethyl substituted with 0-1 R^{5b};
propyl substituted with 0-1 R^{5b}; or
butyl substituted with 0-1 R^{5b};

10

R^{5a} is H;

R^{5b}, at each occurrence, is independently selected from:
H, methyl, ethyl, and NR¹⁵R¹⁶;

15

R¹¹, at each occurrence, is independently selected from
H, NR¹⁸R¹⁹, CF₃;

C₁-C₄ alkyl substituted with 0-1 R^{11a};

phenyl substituted with 0-3 R^{11b};

20

C₃-C₆ carbocycle substituted with 0-3 R^{11b}; or

5 to 7 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle
is substituted with 0-3 R^{11b}; wherein said 5 to 7
membered heterocycle is selected from pyridinyl,
pyrimidinyl, triazinyl, furanyl, thienyl,
thiazolyl, pyrrolyl, piperazinyl, piperidinyl,
homopiperidinyl, pyrazolyl, imidazolyl, oxazolyl,
isoxazolyl, and tetrazolyl;

25

R^{11a}, at each occurrence, is independently selected from H,
methyl, ethyl, propyl, butyl, methoxy, ethoxy,
propoxy, Cl, F, NR¹⁵R¹⁶, CF₃;
phenyl substituted with 0-3 R^{11b};

30

C₃-C₆ carbocycle substituted with 0-3 R^{11b}; or
5 to 7 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle
5 is substituted with 0-3 R^{11b}; wherein said 5 to 7
membered heterocycle is selected from pyridinyl,
pyrimidinyl, triazinyl, furanyl, thienyl,
thiazolyl, pyrrolyl, piperazinyl, piperidinyl,
homopiperidinyl, pyrazolyl, imidazolyl, oxazolyl,
10 isoxazolyl, and tetrazolyl;

R^{11b}, at each occurrence, is independently selected from
H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl,
methoxy, and ethoxy;

15 W is a bond, -CH₂-, or -CH(CH₃)-;

X is a bond;
phenyl substituted with 0-1 R^{Xb};
20 C₃-C₆ cycloalkyl substituted with 0-3 R^{Xb}; or
5 to 6 membered heterocycle containing 1 to 3
heteroatoms selected from nitrogen, oxygen, and
sulphur; wherein said 5 to 6 membered heterocycle
is selected from pyridinyl, pyrimidinyl,
25 triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl,
piperazinyl, piperidinyl, pyrazolyl, imidazolyl,
oxazolyl, and isoxazolyl;

Y is a bond, -V-, -CH₂-V-, or -V-CH₂-;

30 V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-,
-N(R¹⁹)-, -C(=O)NH-, or -NHC(=O)-;

Z is H;
35 C₁-C₆ alkyl substituted with 0-2 R¹²;

- C₂-C₆ alkenyl substituted with 0-2 R¹²;
C₂-C₆ alkynyl substituted with 0-2 R¹²;
phenyl substituted with 0-4 R^{12b};
C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or
5 5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
heterocycle is substituted with 0-3 R^{12b};
- 10 R¹² is phenyl substituted with 0-4 R^{12b};
C₃-C₆ carbocycle substituted with 0-4 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered
15 heterocycle is substituted with 0-3 R^{12b};
- R^{12b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, NR¹⁵R¹⁶, CF₃, OCF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, methyl, ethyl, propyl, butyl,
20 methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂
haloalkoxy;
- R¹³, at each occurrence, is independently selected from
H, methyl, methoxy, Cl, F, Br, and CF₃;
25
- R¹⁴, at each occurrence, is independently selected from H,
phenyl, benzyl, methyl, ethyl, propyl, and butyl;
- R¹⁵, at each occurrence, is independently selected from H,
30 methyl, ethyl, propyl, or butyl;
- R¹⁶, at each occurrence, is independently selected from H,
OH, methyl, ethyl, propyl, butyl, phenyl, benzyl, and
phenethyl;
- 35

R¹⁸, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl;

5 R¹⁹, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, and butyl; and

R²⁰ is H, methyl, ethyl, propyl, or butyl.

10

8. A compound of Claim 6 wherein:

Q is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH₂CH₂CH₂CH₃, -CH(CH₃)₂,
-CH(CH₃)CH₂CH₃, -CH₂CH(CH₃)₂,

15

-CH₂CF₃, -CH₂CH₂CF₃, -CH₂CH₂CH₂CF₃,

phenyl-, 4-tBu-phenyl-, 4-iPr-phenyl-, 4-Et-phenyl-,

2-F-phenyl-, 3-F-phenyl-, 4-F-phenyl-,

20

2-Cl-phenyl-, 3-Cl-phenyl-, 4-Cl-phenyl-,

2-Br-phenyl-, 3-Br-phenyl-, 4-Br-phenyl-,

2-NO₂-phenyl-, 3-NO₂-phenyl-, 4-NO₂-phenyl-,

2-CH₃-phenyl-, 3-CH₃-phenyl-, 4-CH₃-phenyl-,

2-CH₃O-phenyl-, 3-CH₃O-phenyl-, 4-CH₃O-phenyl-,

25

2-CF₃-phenyl-, 3-CF₃-phenyl-, 4-CF₃-phenyl-,

2-CF₃O-phenyl-, 3-CF₃O-phenyl-, 4-CF₃O-phenyl-,

2-CH₃CONH-phenyl, 3-CH₃CONH-phenyl, 4-CH₃CONH-phenyl,

2-methoxycarbonyl-phenyl-, 4-phenyl-phenyl-,

30

2,3-diF-phenyl-, 2,4-diF-phenyl-, 2,5-diF-phenyl-,

2,6-diF-phenyl-, 3,4-diF-phenyl-, 3,5-diF-phenyl-,

2,3-diCl-phenyl-, 2,4-diCl-phenyl-, 2,5-diCl-phenyl-,

2,6-diCl-phenyl-, 3,4-diCl-phenyl-, 3,5-diCl-phenyl-,

3-F-4-Cl-phenyl-, 3-F-5-Cl-phenyl-, 3-Cl-4-F-phenyl-,

35

2,3-diMe-phenyl-, 2,4-diMe-phenyl-, 2,5-diMe-phenyl-,

2,6-diMe-phenyl-, 3,4-diMe-phenyl-, 3,5-diMe-phenyl-,

2,3-diMeO-phenyl-, 2,4-diMeO-phenyl-, 2,5-diMeO-phenyl-,
 2,6-diMeO-phenyl-, 3,4-diMeO-phenyl-, 3,5-diMeO-phenyl-,
 2,3-diCF₃-phenyl-, 2,4-diCF₃-phenyl-, 2,5-diCF₃-phenyl-,
 2,6-diCF₃-phenyl-, 3,4-diCF₃-phenyl-, 3,5-diCF₃-phenyl-,
 5 2-NO₂-4-CF₃-phenyl-, 2-Me-5-Cl-phenyl-,
 2,4,6-triMe-phenyl,

benzyl-, naphth-1-yl-, naphth-2-yl-, beta-styrene-,
 furanyl-, thienyl-, pyridyl-, thiazolyl-, imidazol-1-yl-,
 10 , oxazolyl-, isoxazolyl-, quinolin-8-yl-,

3-methyl-isoxazol-4-yl-,
 3,5-dimethyl-isoxazol-4-yl-,
 3-bromo-5-chloro-thiophen-2-yl-,
 15 2,3-dichlorothiophen-5-yl-,
 4-bromo-5-chlorothiophen-2-yl-,
 5-[(benzoylamino)methyl]-thiophen-2-yl-,
 4-phenylsulfonylthiophen-2-yl-,
 5-(phenylsulfonyl)thiophen-2-yl-,
 20 2-(1-methyl-(5-trifluoromethyl)pyrazole)thiophen-5-yl-,
 5-(2-pyridyl)thiophen-2-yl-,
 1-methyl-5-(trifluoromethyl)imidazol-3-yl-,
 2-(2-methylthio-pyrimidin-3-yl)-thiophen-5-yl-, or
 dibenzofuran-2-yl-;

25 R⁵ is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃,
 -CH(CH₃)CH₂CH₃, -CH₂CH(CH₃)₂, -CH₂C(CH₃)₃,
 -CH₂CH₂CH₂CH₂CH₃, -CH(CH₃)CH₂CH₂CH₃, -CH₂CH(CH₃)CH₂CH₃,
 -CH₂CH₂CH(CH₃)₂, -CH(CH₂CH₃)₂,


30 cyclopropyl-CH₂-, cyclobutyl-CH₂-, cyclopentyl-CH₂-,
 cyclohexyl-CH₂-, (2-CH₃-cyclopropyl)CH₂-,
 (3-CH₃-cyclobutyl)CH₂-, cyclopropyl-CH₂CH₂-,
 cyclobutyl-CH₂CH₂-, cyclopentyl-CH₂CH₂-,
 35 cyclohexyl-CH₂CH₂-, (2-CH₃-cyclopropyl)CH₂CH₂-,
 (3-CH₃-cyclobutyl)CH₂CH₂-,

phenyl-CH₂-, (2-F-phenyl)CH₂-, (3-F-phenyl)CH₂-,
 (4-F-phenyl)CH₂-, furanyl-CH₂-, thienyl-CH₂-,
 pyridyl-CH₂-, 1-imidazolyl-CH₂-, oxazolyl-CH₂-,
 5 isoxazolyl-CH₂-,

phenyl-CH₂CH₂-, (2-F-phenyl)CH₂CH₂-, (3-F-phenyl)CH₂CH₂-
 ,
 (4-F-phenyl)CH₂CH₂-, furanyl-CH₂CH₂-, thienyl-CH₂CH₂-,
 10 pyridyl-CH₂CH₂-, 1-imidazolyl-CH₂CH₂-, oxazolyl-CH₂CH₂-,
 or isoxazolyl-CH₂CH₂-,

W is a bond, -CH₂-, or -CH(CH₃)-;

15 X is a bond;



20

Y is a bond, -CH₂-V-, -V-, or -V-CH₂-;

V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-,
 -N(CH₃)-, -C(=O)NH-, or -NHC(=O)-,

25

Z is H, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂,
 -CH₂CH₂CH₂CH₃, -CH(CH₃)CH₂CH₃, -CH₂CH(CH₃)₂, -C(CH₃)₃,
 -CH₂CH₂CH₂CH₂CH₃, -CH(CH₃)CH₂CH₂CH₃, -CH₂CH(CH₃)CH₂CH₃,
 -CH₂CH₂CH(CH₃)₂, -CH(CH₂CH₃)₂, -CH₂C(CH₃)₃, -
 30 C(CH₃)₂CH₂CH₃,
 -CH(CH₃)CH(CH₃)₂, -CH₂CH₂CH₂CH₂CH₂CH₃,
 -CH(CH₃)CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)CH₂CH₂CH₃,

-CH₂CH₂CH(CH₃)CH₂CH₃, -CH₂CH₂CH₂CH(CH₃)₂, -
 CH₂CH(CH₂CH₃)₂,
 -CH₂CH₂C(CH₃)₃, -CH(CH₂CH₃)CH₂CH₂CH₃,
 -CF₃, -CH₂CF₃, -CH₂CH₂CF₃, -CH₂CH₂CH₂CF₃, -
 5 CH₂CH₂CH₂CH₂CF₃,
 cyclopropyl-, (cyclopropyl)CH₂-, (cyclopropyl)CH₂CH₂-,
 cyclobutyl-, (cyclobutyl)CH₂-, (cyclobutyl)CH₂CH₂-,
 cyclopentyl-, (cyclopentyl)CH₂-, (cyclopentyl)CH₂CH₂-,
 cyclohexyl-, (cyclohexyl)CH₂-, (cyclohexyl)CH₂CH₂-,
 10 phenyl, 2-F-phenyl, 3-F-phenyl, 4-F-phenyl, 2-Cl-phenyl,
 3-Cl-phenyl, 4-Cl-phenyl, 2,3-diF-phenyl,
 2,4-diF-phenyl, 2,5-diF-phenyl, 2,6-diF-phenyl,
 3,4-diF-phenyl, 3,5-diF-phenyl, 2,3-diCl-phenyl,
 2,4-diCl-phenyl, 2,5-diCl-phenyl, 2,6-diCl-phenyl,
 15 3,4-diCl-phenyl, 3,5-diCl-phenyl, 3-F-4-Cl-phenyl,
 3-F-5-Cl-phenyl, 3-Cl-4-F-phenyl, 2-MeO-phenyl,
 3-MeO-phenyl, 4-MeO-phenyl, 2-Me-phenyl, 3-Me-phenyl,
 4-Me-phenyl, 2-MeS-phenyl, 3-MeS-phenyl, 4-MeS-phenyl,
 2-CF₃O-phenyl, 3-CF₃O-phenyl, 4-CF₃O-phenyl, furanyl,
 20 thienyl, pyridyl, 2-Me-pyridyl, 3-Me-pyridyl,
 4-Me-pyridyl, 1-imidazolyl, oxazolyl, isoxazolyl,
 1-benzimidazolyl, morpholino, N-piperidinyl,
 phenyl-CH₂-, (2-F-phenyl)CH₂-, (3-F-phenyl)CH₂-,
 (4-F-phenyl)CH₂-, (2-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂-,
 25 (4-Cl-phenyl)CH₂-, (2,3-diF-phenyl)CH₂-,
 (2,4-diF-phenyl)CH₂-, (2,5-diF-phenyl)CH₂-,
 (2,6-diF-phenyl)CH₂-, (3,4-diF-phenyl)CH₂-,
 (3,5-diF-phenyl)CH₂-, (2,3-diCl-phenyl)CH₂-,
 (2,4-diCl-phenyl)CH₂-, (2,5-diCl-phenyl)CH₂-,
 30 (2,6-diCl-phenyl)CH₂-, (3,4-diCl-phenyl)CH₂-,
 (3,5-diCl-phenyl)CH₂-, (3-F-4-Cl-phenyl)CH₂-,
 (3-F-5-Cl-phenyl)CH₂-, (3-Cl-4-F-phenyl)CH₂-,
 (2-MeO-phenyl)CH₂-, (3-MeO-phenyl)CH₂-,
 (4-MeO-phenyl)CH₂-, (2-Me-phenyl)CH₂-,

(3-Me-phenyl)CH₂-, (4-Me-phenyl)CH₂-, (2-MeS-phenyl)CH₂-
, (3-MeS-phenyl)CH₂-, (4-MeS-phenyl)CH₂-,
(2-CF₃O-phenyl)CH₂-, (3-CF₃O-phenyl)CH₂-,
(4-CF₃O-phenyl)CH₂-, (furanyl)CH₂-, (thienyl)CH₂-,
5 (pyridyl)CH₂-, (2-Me-pyridyl)CH₂-, (3-Me-pyridyl)CH₂-,
(4-Me-pyridyl)CH₂-, (1-imidazolyl)CH₂-, (oxazolyl)CH₂-,
(isoxazolyl)CH₂-, (1-benzimidazolyl)CH₂-,
morpholino)CH₂-, (N-piperidiny)CH₂-,
phenyl-CH₂CH₂-, (phenyl)₂CHCH₂-, (2-F-phenyl)CH₂CH₂-,
10 (3-F-phenyl)CH₂CH₂-, (4-F-phenyl)CH₂CH₂-,
(2-Cl-phenyl)CH₂CH₂-, (3-Cl-phenyl)CH₂CH₂-,
(4-Cl-phenyl)CH₂CH₂-, (2,3-diF-phenyl)CH₂CH₂-,
(2,4-diF-phenyl)CH₂CH₂-, (2,5-diF-phenyl)CH₂CH₂-,
(2,6-diF-phenyl)CH₂CH₂-, (3,4-diF-phenyl)CH₂CH₂-,
15 (3,5-diF-phenyl)CH₂CH₂-, (2,3-diCl-phenyl)CH₂CH₂-,
(2,4-diCl-phenyl)CH₂CH₂-, (2,5-diCl-phenyl)CH₂CH₂-,
(2,6-diCl-phenyl)CH₂CH₂-, (3,4-diCl-phenyl)CH₂CH₂-,
(3,5-diCl-phenyl)CH₂CH₂-, (3-F-4-Cl-phenyl)CH₂CH₂-,
(3-F-5-Cl-phenyl)CH₂CH₂-, (3-Cl-4-F-phenyl)CH₂CH₂-,
20 (2-MeO-phenyl)CH₂CH₂-, (3-MeO-phenyl)CH₂CH₂-,
(4-MeO-phenyl)CH₂CH₂-, (2-Me-phenyl)CH₂CH₂-,
(3-Me-phenyl)CH₂CH₂-, (4-Me-phenyl)CH₂CH₂-,
(2-MeS-phenyl)CH₂CH₂-, (3-MeS-phenyl)CH₂CH₂-,
(4-MeS-phenyl)CH₂CH₂-, (2-CF₃O-phenyl)CH₂CH₂-,
25 (3-CF₃O-phenyl)CH₂CH₂-, (4-CF₃O-phenyl)CH₂CH₂-,
(furanyl)CH₂CH₂-, (thienyl)CH₂CH₂-, (pyridyl)CH₂CH₂-,
(2-Me-pyridyl)CH₂CH₂-, (3-Me-pyridyl)CH₂CH₂-,
(4-Me-pyridyl)CH₂CH₂-, (imidazolyl)CH₂CH₂-,
(oxazolyl)CH₂CH₂-, (isoxazolyl)CH₂CH₂-,
30 (benzimidazolyl)CH₂CH₂-, (cyclopropyl)CH₂CH₂-,
(cyclobutyl)CH₂CH₂-, (cyclopentyl)CH₂CH₂-,
(cyclohexyl)CH₂CH₂-, (morpholino)CH₂CH₂-, or
(N-piperidiny)CH₂CH₂-;

- R¹¹, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, cyclopropylethyl, cyclobutylethyl, cyclopentylethyl, cyclohexylethyl, 4-F-phenyl, 4-Cl-phenyl, 4-CH₃-phenyl, 4-CF₃-phenyl, 4-CH₃O-phenyl, 4-CF₃O-phenyl, 3-F-phenyl, 3-Cl-phenyl, 3-CH₃-phenyl, 3-CF₃-phenyl, 3-CH₃O-phenyl, 3-CF₃O-phenyl, 2-F-phenyl, 2-Cl-phenyl, 2-CH₃-phenyl, 2-CF₃-phenyl, 2-CH₃O-phenyl, 2-CF₃O-phenyl, (4-F-phenyl)methyl-, (4-Cl-phenyl)methyl-, (4-CH₃-phenyl)methyl-, (4-CF₃-phenyl)methyl-, (4-CH₃O-phenyl)methyl-, (4-CF₃O-phenyl)methyl-, (3-F-phenyl)methyl-, (3-Cl-phenyl)methyl-, (3-CH₃-phenyl)methyl-, (3-CF₃-phenyl)methyl-, (3-CH₃O-phenyl)methyl-, (3-CF₃O-phenyl)methyl-, (2-F-phenyl)methyl-, (2-Cl-phenyl)methyl-, (2-CH₃-phenyl)methyl-, (2-CF₃-phenyl)methyl-, (2-CH₃O-phenyl)methyl-, (2-CF₃O-phenyl)methyl-, 2-pyridyl-, 3-pyridyl-, 4-pyridyl-, 1-piperidiny, 1-homopiperidiny, and 1-morpholino; and
- R¹³, at each occurrence, is independently selected from H, F, Cl, and methoxy.

9. A compound of Claim 7 wherein:

- Q is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, phenyl-, 2-F-phenyl-, 2-Cl-phenyl-, 2-Br-phenyl-, 2-NO₂-phenyl-, 2-CH₃-phenyl-, 2-CH₃CH₂-phenyl-, 2-CH₃O-phenyl-, 2-CF₃-phenyl-, 2-CF₃O-phenyl-, 2-CH₃CONH-phenyl, or 3,5-dimethyl-isoxazol-4-yl-;


R^5 is $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$,
 $-CH(CH_3)CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH_2C(CH_3)_3$,
 $-CH_2CH_2CH_2CH_2CH_3$, $-CH(CH_3)CH_2CH_2CH_3$, $-CH_2CH(CH_3)CH_2CH_3$,
 $-CH_2CH_2CH(CH_3)_2$, $-CH(CH_2CH_3)_2$,

5

W is a bond, $-CH_2-$, or $-CH(CH_3)-$;

X is a bond;

10



Y is a bond, $-CH_2-V-$, $-V-$, or $-V-CH_2-$;

15

V is a bond, $-C(=O)-$, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$,
 $-N(CH_3)-$, $-C(=O)NH-$, or $-NH\dot{C}(=O)-$,

Z is H, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$,

20

$-CH_2CH_2CH_2CH_3$, $-CH(CH_3)CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-C(CH_3)_3$,
 $-CH_2CH_2CH_2CH_2CH_3$, $-CH(CH_3)CH_2CH_2CH_3$, $-CH_2CH(CH_3)CH_2CH_3$,
 $-CH_2CH_2CH(CH_3)_2$, $-CH(CH_2CH_3)_2$, $-CH_2C(CH_3)_3$, $-$
 $C(CH_3)_2CH_2CH_3$,

25

$-CH(CH_3)CH(CH_3)_2$, $-CH_2CH_2CH_2CH_2CH_2CH_3$,
 $-CH(CH_3)CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)CH_2CH_2CH_3$,
 $-CH_2CH_2CH(CH_3)CH_2CH_3$, $-CH_2CH_2CH_2CH(CH_3)_2$, $-$
 $CH_2CH(CH_2CH_3)_2$,
 $-CH_2CH_2C(CH_3)_3$, $-CH(CH_2CH_3)CH_2CH_2CH_3$,
 $-CF_3$, $-CH_2CF_3$, $-CH_2CH_2CF_3$, $-CH_2CH_2CH_2CF_3$, $-$

30

$CH_2CH_2CH_2CH_2CF_3$,
cyclopropyl-, (cyclopropyl) CH_2- , (cyclopropyl) CH_2CH_2- ,
cyclobutyl-, (cyclobutyl) CH_2- , (cyclobutyl) CH_2CH_2- ,

cyclopentyl-, (cyclopentyl)CH₂-, (cyclopentyl)CH₂CH₂-,
cyclohexyl-, (cyclohexyl)CH₂-, (cyclohexyl)CH₂CH₂-,
phenyl, 2-F-phenyl, 3-F-phenyl, 4-F-phenyl, 2-Cl-phenyl,
3-Cl-phenyl, 4-Cl-phenyl, 2,3-diF-phenyl,
5 2,4-diF-phenyl, 2,5-diF-phenyl, 2,6-diF-phenyl,
3,4-diF-phenyl, 3,5-diF-phenyl, 2,3-diCl-phenyl,
2,4-diCl-phenyl, 2,5-diCl-phenyl, 2,6-diCl-phenyl,
3,4-diCl-phenyl, 3,5-diCl-phenyl, 3-F-4-Cl-phenyl,
3-F-5-Cl-phenyl, 3-Cl-4-F-phenyl, 2-MeO-phenyl,
10 3-MeO-phenyl, 4-MeO-phenyl, 2-Me-phenyl, 3-Me-phenyl,
4-Me-phenyl, 2-MeS-phenyl, 3-MeS-phenyl, 4-MeS-phenyl,
2-CF₃O-phenyl, 3-CF₃O-phenyl, 4-CF₃O-phenyl, furanyl,
thienyl, pyridyl, 2-Me-pyridyl, 3-Me-pyridyl,
4-Me-pyridyl, 1-imidazolyl, oxazolyl, isoxazolyl,
15 1-benzimidazolyl, morpholino, N-piperidinyl,
phenyl-CH₂-, (2-F-phenyl)CH₂-, (3-F-phenyl)CH₂-,
(4-F-phenyl)CH₂-, (2-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂-,
(4-Cl-phenyl)CH₂-, (2,3-diF-phenyl)CH₂-,
(2,4-diF-phenyl)CH₂-, (2,5-diF-phenyl)CH₂-,
20 (2,6-diF-phenyl)CH₂-, (3,4-diF-phenyl)CH₂-,
(3,5-diF-phenyl)CH₂-, (2,3-diCl-phenyl)CH₂-,
(2,4-diCl-phenyl)CH₂-, (2,5-diCl-phenyl)CH₂-,
(2,6-diCl-phenyl)CH₂-, (3,4-diCl-phenyl)CH₂-,
(3,5-diCl-phenyl)CH₂-, (3-F-4-Cl-phenyl)CH₂-,
25 (3-F-5-Cl-phenyl)CH₂-, (3-Cl-4-F-phenyl)CH₂-,
(2-MeO-phenyl)CH₂-, (3-MeO-phenyl)CH₂-,
(4-MeO-phenyl)CH₂-, (2-Me-phenyl)CH₂-,
(3-Me-phenyl)CH₂-, (4-Me-phenyl)CH₂-, (2-MeS-phenyl)CH₂-,
(3-MeS-phenyl)CH₂-, (4-MeS-phenyl)CH₂-,
30 (2-CF₃O-phenyl)CH₂-, (3-CF₃O-phenyl)CH₂-,
(4-CF₃O-phenyl)CH₂-, (furanyl)CH₂-, (thienyl)CH₂-,
(pyridyl)CH₂-, (2-Me-pyridyl)CH₂-, (3-Me-pyridyl)CH₂-,
(4-Me-pyridyl)CH₂-, (1-imidazolyl)CH₂-, (oxazolyl)CH₂-,
(isoxazolyl)CH₂-, (1-benzimidazolyl)CH₂-,
35 morpholino)CH₂-, (N-piperidinyl)CH₂-,


phenyl-CH₂CH₂-, (phenyl)₂CHCH₂-, (2-F-phenyl)CH₂CH₂-,
 (3-F-phenyl)CH₂CH₂-, (4-F-phenyl)CH₂CH₂-,
 (2-Cl-phenyl)CH₂CH₂-, (3-Cl-phenyl)CH₂CH₂-,
 (4-Cl-phenyl)CH₂CH₂-, (2,3-diF-phenyl)CH₂CH₂-,
 5 (2,4-diF-phenyl)CH₂CH₂-, (2,5-diF-phenyl)CH₂CH₂-,
 (2,6-diF-phenyl)CH₂CH₂-, (3,4-diF-phenyl)CH₂CH₂-,
 (3,5-diF-phenyl)CH₂CH₂-, (2,3-diCl-phenyl)CH₂CH₂-,
 (2,4-diCl-phenyl)CH₂CH₂-, (2,5-diCl-phenyl)CH₂CH₂-,
 (2,6-diCl-phenyl)CH₂CH₂-, (3,4-diCl-phenyl)CH₂CH₂-,
 10 (3,5-diCl-phenyl)CH₂CH₂-, (3-F-4-Cl-phenyl)CH₂CH₂-,
 (3-F-5-Cl-phenyl)CH₂CH₂-, (3-Cl-4-F-phenyl)CH₂CH₂-,
 (2-MeO-phenyl)CH₂CH₂-, (3-MeO-phenyl)CH₂CH₂-,
 (4-MeO-phenyl)CH₂CH₂-, (2-Me-phenyl)CH₂CH₂-,
 (3-Me-phenyl)CH₂CH₂-, (4-Me-phenyl)CH₂CH₂-,
 15 (2-MeS-phenyl)CH₂CH₂-, (3-MeS-phenyl)CH₂CH₂-,
 (4-MeS-phenyl)CH₂CH₂-, (2-CF₃O-phenyl)CH₂CH₂-,
 (3-CF₃O-phenyl)CH₂CH₂-, (4-CF₃O-phenyl)CH₂CH₂-,
 (furanyl)CH₂CH₂-, (thienyl)CH₂CH₂-, (pyridyl)CH₂CH₂-,
 (2-Me-pyridyl)CH₂CH₂-, (3-Me-pyridyl)CH₂CH₂-,
 20 (4-Me-pyridyl)CH₂CH₂-, (imidazolyl)CH₂CH₂-,
 (oxazolyl)CH₂CH₂-, (isoxazolyl)CH₂CH₂-,
 (benzimidazolyl)CH₂CH₂-, (cyclopropyl)CH₂CH₂-,
 (cyclobutyl)CH₂CH₂-, (cyclopentyl)CH₂CH₂-,
 (cyclohexyl)CH₂CH₂-, (morpholino)CH₂CH₂-, or
 25 (N-piperidiny)CH₂CH₂-;

R¹¹, at each occurrence, is independently selected from H,
 methyl, ethyl, propyl, butyl, phenyl, benzyl, phenethyl,
 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,
 30 cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl,
 cyclohexylmethyl, cyclopropylethyl, cyclobutylethyl,
 cyclopentylethyl, cyclohexylethyl, 4-F-phenyl,
 4-Cl-phenyl, 4-CH₃-phenyl, 4-CF₃-phenyl, 4-CH₃O-phenyl,
 4-CF₃O-phenyl, 3-F-phenyl, 3-Cl-phenyl, 3-CH₃-phenyl,

3-CF₃-phenyl, 3-CH₃O-phenyl, 3-CF₃O-phenyl, 2-F-phenyl,
2-Cl-phenyl, 2-CH₃-phenyl, 2-CF₃-phenyl, 2-CH₃O-phenyl,
2-CF₃O-phenyl, (4-F-phenyl)methyl-, (4-Cl-phenyl)methyl-

- 5 (4-CH₃-phenyl)methyl-, (4-CF₃-phenyl)methyl-,
(4-CH₃O-phenyl)methyl-, (4-CF₃O-phenyl)methyl-,
(3-F-phenyl)methyl-, (3-Cl-phenyl)methyl-,
(3-CH₃-phenyl)methyl-, (3-CF₃-phenyl)methyl-,
(3-CH₃O-phenyl)methyl-, (3-CF₃O-phenyl)methyl-,
10 (2-F-phenyl)methyl-, (2-Cl-phenyl)methyl-,
(2-CH₃-phenyl)methyl-, (2-CF₃-phenyl)methyl-,
(2-CH₃O-phenyl)methyl-, (2-CF₃O-phenyl)methyl-,
2-pyridyl-, 3-pyridyl-, 4-pyridyl-,
1-piperidinyl, 1-homopiperidinyl, and 1-morpholino; and
15 R¹³, at each occurrence, is independently selected from H,
F, Cl, and methoxy.

- 20 10. A compound, according to Claim 4, 5, 6, 7, 8, or
9, of Formula (Ic);




(Ic)

- 25 or a pharmaceutically acceptable salt form or prodrug
thereof.


11. A compound, according to Claim 4, 5, 6, 7, 8, or 9, of
Formula (Id);

30



or a pharmaceutically acceptable salt form or prodrug
5 thereof.

12. A compound, according to Claim 4, 5, 6, 7, 8, or 9, of
Formula (Ia):



or a pharmaceutically acceptable salt or prodrug thereof.

13. A compound of Claim 1, 2, 3, 4, 5, or 6 wherein R⁵ is
15 OR¹⁴.

14. A compound of Claim 1, 2, 3, or 4 wherein R⁵ is C₁-C₄
alkoxy substituted with 0-3 R^{5b}.

20 15. A compound selected from:

2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

25

2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-(1-
methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-
3-yl)-propionamide;

2-(3-bromo-5-chloro-thiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

5 2-(naphthalene-1-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

10 2-(5-dimethylamino naphthalene-1-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

15 2-(naphthalene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

20 2-(2-acetamido-4-methylthiazole-5-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

2-(thiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

25 2-(quinoline-8-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

30 2-(phenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

35 2-(2,5-dichlorophenyl sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

2-(mesitylene-2-sulfonylamino)-4-methyl-pentanoic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

5 2-(3-nitrophenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

10 2-(4-bromophenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

15 2-(4-fluorophenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

20 2-(4-chlorophenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

2-(4-acetamidophenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

25 2-(4-nitrophenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

30 2-(4-methoxyphenyl-sulfonylamino)-4-methyl-pentanoic
acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

35 2-(4-tert-butylphenyl-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

2-(p-toluene-sulfonylamino)-4-methyl-pentanoic acid
(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

5 2-(benzyl-sulfonylamino)-4-methyl-pentanoic acid (1-
methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-
3-yl)-amide;

2-(beta-styrene-sulfonylamino)-4-methyl-pentanoic acid
10 (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

2-((2-methoxycarbonyl)phenyl-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
15 benzo[e][1,4]diazepin-3-yl)-amide;

2-(2-nitro-4-(trifluoromethyl)phenyl-sulfonylamino)-4-
methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-
1H-benzo[e][1,4]diazepin-3-yl)-amide;

20 2-(3-(trifluoromethyl)phenyl-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

25 2-(2,5-dimethoxyphenyl-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

2-(2-methylphenyl-sulfonylamino)-4-methyl-pentanoic
30 acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-amide;

2-(3,4-dichlorophenyl-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
35 benzo[e][1,4]diazepin-3-yl)-amide;

2-(4-(trifluoromethoxy)phenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

5 2-(3,4-dimethoxyphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

10 2-(2-bromophenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

15 2-(3,5-bis(trifluoromethyl)phenyl-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

20 2-(4-ethylphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

2-(4-isopropylphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

25 2-(2,5-dichlorothiophene-3-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

30 2-(5-chlorothiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

35 2-(2-(trifluoromethyl)phenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

2-(3-methylphenyl-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

5 2-(2,3-dichlorothiophene-5-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

2-(4-bromo-5-chlorothiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

10

2-(5-[(benzoylamino)methyl]-thiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

15

2-(4-phenylsulfonylthiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

20

2-(5-(phenylsulfonyl)thiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

25 2-[2-(1-methyl-(5-trifluoromethyl)pyrazole)thiophene-5-sulfonylamino]-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

2-(5-(2-pyridyl)thiophene-2-sulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

30

2-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-4-methyl-pentanoic acid [5-(4-fluoro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-amide;

35

N-[1-(3'-Chloro-4'-fluoro-biphenyl-3-ylmethyl)-2-oxo-azepan-3-yl]-2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-propionamide;

5 2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-[5-(4-fluoro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-propionamide;

10 2-(2-fluoro-benzenesulfonylamino)-N-[5-(4-fluoro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-propionamide;

15 N-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-2-(2-trifluoromethyl-benzenesulfonylamino)-propionamide;

20 2-(2-fluoro-benzenesulfonylamino)-4-methyl-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

20 4-methyl-2-(2-trifluoromethyl-benzenesulfonylamino)-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

25 4-methyl-2-(propane-1-sulfonylamino)-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

30 4-methyl-2-(propane-1-sulfonylamino)-pentanoic acid (1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

35 2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-N-[1-methyl-2-oxo-5-(4-trifluoromethyl-phenyl)-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl]-propionamide;

2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-4-methyl-pentanoic acid (7-chloro-1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-amide;

4-methyl-2-(propane-1-sulfonylamino)-pentanoic acid
[5-(4-chloro-phenyl)-1-methyl-2-oxo-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl]-amide;

5

2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-4-methyl-
pentanoic acid (1-methyl-2-oxo-5-pyridin-2-yl-2,3-dihydro-
1H-benzo[e][1,4]diazepin-3-yl)-amide;

10

2-[(3,5-dimethyl-isoxazole-4-sulfonyl)-methyl-amino]-
N-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-propionamide;

15

3-dimethylamino-2-(3,5-dimethyl-isoxazole-4-
sulfonylamino)-N-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-propionamide;

20

N-[1-methyl-2-oxo-5-(4-trifluoromethyl-phenyl)-2,3-
dihydro-1H-benzo[e][1,4]diazepin-3-yl]-2-(propane-1-
sulfonylamino)-propionamide;

25

N-(7-chloro-1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl)-2-(3,5-dimethyl-isoxazole-4-
sulfonylamino)-propionamide;

2-(3,5-dimethyl-isoxazole-4-sulfonylamino)-4-methyl-
pentanoic acid [5-(4-chloro-phenyl)-1-methyl-2-oxo-2,3-
dihydro-1H-benzo[e][1,4]diazepin-3-yl]-amide; and

30

2-methanesulfonylamino-N-[1-methyl-2-oxo-5-(4-
trifluoromethyl-phenyl)-2,3-dihydro-1H-
benzo[e][1,4]diazepin-3-yl]-propionamide.

35

16. A pharmaceutical composition comprising a compound of
Claim 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 and a
pharmaceutically acceptable carrier.

17. A method for the treatment of neurological disorders

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 00/27666

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D413/12 A61K31/55 C07D409/12 C07D401/12 C07D409/14
 C07D403/12 C07D407/12 C07D243/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 99 67221 A (MCDANIEL STACEY L ;AUDIA JAMES E (US); CWI CYNTHIA L (US); ELAN PH) 29 December 1999 (1999-12-29) page 97; table 6.1 page 401, line 25 -page 404, line 25; claims 1-123 ---	1-21
X	US 5 852 007 A (CHATTERJEE SANKAR) 22 December 1998 (1998-12-22) column 64; claim 1; example 108 column 1, line 21 -column 2, line 22 ---	1-21
P,A	WO 00 07995 A (DU PONT PHARM CO) 17 February 2000 (2000-02-17) the whole document --- -/--	1-21



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

30 November 2000

Date of mailing of the international search report

19 12 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Schmid, J-C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/27666

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99 32453 A (MCDANIEL STACEY L ;AUDIA JAMES E (US); ELAN PHARM INC (US); LILLY) 1 July 1999 (1999-07-01) the whole document -----	1-21

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 00/27666**Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)**

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 17-21 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern al Application No

PCT/US 00/27666

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9967221 A	29-12-1999	AU 4707999 A	10-01-2000
		AU 4710199 A	10-01-2000
		AU 4710499 A	10-01-2000
		WO 9967219 A	29-12-1999
		WO 9966934 A	29-12-1999
US 5852007 A	22-12-1998	AU 714324 B	06-01-2000
		AU 1025397 A	03-07-1997
		CA 2238175 A	19-06-1997
		EP 0910564 A	28-04-1999
		WO 9721690 A	19-06-1997
WO 0007995 A	17-02-2000	AU 5337899 A	28-02-2000
WO 9932453 A	01-07-1999	AU 1277799 A	12-07-1999
		BR 9812773 A	10-10-2000
		EP 1042298 A	11-10-2000